

## Evaluation and Treatment of Dyspareunia

John F. Steege, MD, and Denniz A. Zolnoun, MD, MPH  
*Obstet Gynecol* 2009;113(5)

---

### ACCME Accreditation

The American College of Obstetricians and Gynecologists (ACOG) is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians. (*Continuing medical education credit for "Evaluation and Treatment of Dyspareunia" will be available through May 2012.*)

### AMA PRA Category 1 Credit™ and ACOG Cognate Credit

The American College of Obstetricians and Gynecologists (ACOG) designates this educational activity for a maximum of 2 AMA PRA Category 1 Credits™ or up to a maximum of 2 Category 1 ACOG cognate credits. Physicians should only claim credit commensurate with the extent of their participation in the activity.

### Disclosure Statement

Current guidelines state that continuing medical education (CME) providers must ensure that CME activities are free from the control of any commercial interest. All authors, reviewers, and contributors have disclosed to ACOG all relevant financial relationships with any commercial interests. The authors, reviewers, and contributors declare that neither they nor any business associate nor any member of their immediate families has financial interest or other relationships with any manufacturer of products or any providers of services discussed in this program. Any conflicts have been resolved through group and outside review of all content.

Before submitting this form, please print a completed copy as confirmation of your program participation.

*ACOG Fellows:* To obtain credits, complete and return this form by clicking on "Submit" at the bottom of the page. Credit will be automatically recorded upon receipt and online transcripts will be updated twice monthly. ACOG Fellows may check their transcripts online at <http://www.acog.org>.

*Non-ACOG Fellows:* To obtain credits, submit the printout of the completed quiz to your accrediting institution. The printout of the completed quiz is documentation for your continuing medical education credits.

---

1. The embryologic anlage of the vaginal vestibule includes the:

- A. Ovarian parenchyma
- B. Fallopian tube
- C. Cervix and uterus
- D. Rectum and anal verge
- E. Urethra and bladder

2. The main nerve supply to the vulva is via the:
  - A. Pudendal nerve
  - B. Iliohypogastric nerve
  - C. Anococcygeal nerve
  - D. Genitofemoral nerve
  - E. Ilioinguinal nerve
  
3. The concept of neuroplasticity suggests that painful sensations may occur from:
  - A. Altered central signaling pathways
  - B. New peripheral neural connections
  - C. Peripheral physiologic changes
  - D. Repeated subthreshold negative stimuli
  - E. Central desensitization
  
4. Vaginal lubrication in response to sexual stimulation is derived from the:
  - A. Endocervical glands
  - B. Vaginal epithelium
  - C. Bartholin's gland
  - D. Skene's gland
  - E. All of the above
  
5. Myofascial abdominal wall components of dyspareunia are best demonstrated by:
  - A. Palpation of the abdominal wall with a single finger
  - B. The application of ice to the abdominal wall
  - C. The response to muscle relaxant medications
  - D. Having the patient lift her head during examination
  - E. Bimanual pelvic examination

6. In addition to the vagina, lichen planus can also affect the:

- A. Gingiva
- B. Rectum
- C. Extensor surfaces of the skin
- D. Nipples
- E. Bartholin's gland

7. Vulvar vestibulitis is currently classified as what type of disorder?

- A. Inflammatory
- B. Depressive
- C. Neuropsychiatric
- D. Autoimmune
- E. Neurosensory

8. Overall, approximately what percentage of women have dyspareunia at 6 months after delivery?

- A. 10%
- B. 25%
- C. 50%
- D. 75%
- E. 90%

9. Retroversion of the uterus is present in approximately what percent of normal women?

- A. 5%
- B. 10%
- C. 20%
- D. 30%
- E. 40%

10. When a surgical revision of the vaginal apex for deep dyspareunia following hysterectomy is performed, approximately what percentage of patients will have recurrent pain within 2–3 years of surgery?

- A. 5%
- B. 10%
- C. 30%
- D. 50%
- E. 70%

**ACOG ID Number:**

**Name:**

**Address:**

**City/State/Zip:**

**Actual time spent completing this activity (you may record up to 2 hours):**