

TABLE 2. Age and sex distribution, and the location of tumor for the different pathologic diagnoses (n=38)

Diagnosis	Average age \pm SD at presentation (yrs)	Average age \pm SD at exenteration (yrs)	Sex distribution	Location (MC/LC = medial/lateral canthus)
Ocular surface SCC (n=9)	67.7 \pm 12.8	68.1 \pm 12.6	6 men, 3 women	
Periocular SCC (n=6)	56.7 \pm 17.1	57.3 \pm 17.8	6 men	2 MC, 2 LC, 2 lower lid
BCC (n=10)	68.5 \pm 14.7	72.4 \pm 11.5	8 men, 2 women	6 MC
MM (n=5)	76.5 \pm 9.7	81.5 \pm 8	2 men, 3 women	3 uveal, 2 conjunctival
SGC (n=5)	73 \pm 3.2	73.8 \pm 2.2	3 men, 2 women	4 eyelids, 1 conjunctival
Pleomorphic adenocarcinoma (n=1)	48	48	1 woman	Lacrimal gland
Orbital infections (n=2)	75.5 \pm 3.5	76.5 \pm 4.9	1 man, 1 woman	1 MRSA socket infection, 1 mucormycosis

BCC, Basal cell carcinoma; SCC, Squamous cell carcinoma; SGC, Sebaceous gland carcinoma; MM, Malignant melanoma; MRSA, Methicillin-Resistant Staphylococcus aureus