

Practice Guidelines for Perioperative Blood Transfusion and Adjuvant Therapies
American Society of Anesthesiologists

Bibliography

1. Alanay A, Acaroglu E, Ozdemir O, Ercelen O, Bulutcu E, Surat A: Effects of deamino-8-D-arginin vasopressin on blood loss and coagulation factors in scoliosis surgery. A double-blind randomized clinical trial. *Spine* 1999; 24:877-882
2. Alderman EL, Levy JH, Rich JB, Nili M, Vidne B, Schaff H, Uretzky G, Pettersson G, This JJ, Hantler CB, Chaitman B, Nadel A: Analyses of coronary graft patency after aprotinin use: results from the International Multicenter Aprotinin Graft Patency Experience (IMAGE) trial. *J Thorac Cardiovasc Surg* 1998; 116:716-730
3. Almeida MF: Preoperative normovolemic hemodilution in aesthetic plastic surgery. *Aesthetic Plast Surg* 1999; 23:445-449
4. Alvarez JM, Jackson LR, Chatwin C, Smolich JJ: Low-dose postoperative aprotinin reduces mediastinal drainage and blood product use in patients undergoing primary coronary artery bypass grafting who are taking aspirin: a prospective, randomized, double-blind, placebo-controlled trial. *J Thorac Cardiovasc Surg* 2001; 122:457-463
5. Alvarez JM, Quiney NF, McMillan D, Joscelyne K, Connelly T, Brady P, Deal C, Wilson R: The use of ultra-low-dose aprotinin to reduce blood loss in cardiac surgery. *J Cardiothorac Vasc Anesth* 1995; 9:29-33
6. Amar D, Grant FM, Zhang H, Boland PJ, Leung DH, Healey JA: Antifibrinolytic therapy and perioperative blood loss in cancer patients undergoing major orthopedic surgery. *Anesthesiology* 2003; 98:337-342
7. Andrews CM, Lane DW, Bradley JG: Iron pre-load for major joint replacement. *Transfus Med* 1997; 7:281-286
8. Armellin G, Sorbara C, Bonato R, Pittarello D, Cero PD, Giron G: Intraoperative plasmapheresis in cardiac surgery. *J Cardiothorac Vasc Anesth* 1997; 11:13-17
9. Avall A, Hyllner M, Bengtson JP, Carlsson L, Bengtsson A: Greater increase in cytokine concentration after salvage with filtered whole blood than with washed red cells, but no difference in postoperative hemoglobin recovery. *Transfusion* 1999; 39:271-276
10. Avall A, Hyllner M, Bengtson JP, Carlsson L, Bengtsson A: Postoperative inflammatory response after autologous and allogeneic blood transfusion. *Anesthesiology* 1997; 87:511-516
11. Avall A, Hyllner M, Swolin B, Bengtson J, Carlsson L, Bengtsson A: Increased serum erythropoietin concentration after allogeneic compared with autologous blood transfusion. *Transfus Apheresis Sci* 2002; 27:203-210
12. Axford TC, Dearani JA, Ragno G, MacGregor H, Patel MA, Valeri CR, Khuri SF: Safety and therapeutic effectiveness of reinfused shed blood after open heart surgery. *Ann Thorac Surg* 1994; 57:615-622
13. Bacher A, Mayer N, Rajek AM, Haider W: Acute normovolaemic haemodilution does not aggravate gastric mucosal acidosis during cardiac surgery. *Intensive Care Med* 1998; 24:313-321
14. Bailey CR, Kelleher AA, Wielogorski AK: Randomized placebo-controlled double-blind study of three aprotinin regimens in primary cardiac surgery. *Br J Surg* 1994; 81:969-973
15. Barnette RE, Wendling WW, Schweiger JW, Brister NW, Schartel SA, Chen D, Shuman CA, McClurken JB, Jeevanandam V: Intravenous vitamin K1 prior to orthotopic heart transplantation: effects in vivo and in vitro. *Acta Anaesthesiol Scand* 1997; 41:78-83

16. Bartels C, Bechtel JV, Winkler C, Horsch S: Intraoperative autotransfusion in aortic surgery: comparison of whole blood autotransfusion versus cell separation. *J Vasc Surg* 1996; 24:102-108
17. Bartels C, Claeys L, Ktenidis K, Nigbur H, Horsch S: Intraoperative whole blood autotransfusion during venous thrombectomy. *J Cardiovasc Surg* 1994; 35:109-114
18. Basora M, Gomar C, Escolar G, Pacheco M, Fita G, Rodriguez E, Ordinas A: Platelet function during cardiac surgery and cardiopulmonary bypass with low-dose aprotinin. *J Cardiothorac Vasc Anesth* 1999; 13:382-387
19. Beath SM, Nuttall GA, Fass DN, Oliver WC Jr, Ereth MH, Oyen LJ: Plasma aprotinin concentrations during cardiac surgery: full- versus half-dose regimens. *Anesth Analg* 2000; 91:257-264
20. Bedirhan MA, Turna A, Yagan N, Tasci O: Aprotinin reduces postoperative bleeding and the need for blood products in thoracic surgery: results of a randomized double-blind study. *Eur J Cardiothorac Surg* 2001; 20:1122-1127
21. Benli IT, Akalin S, Duman E, Citak M, Kis M: The results of intraoperative autotransfusion in orthopaedic surgery. *Bull Hosp Jt Dis* 1999; 58:184-187
22. Bennett SR: Perioperative autologous blood transfusion in elective total hip prosthesis operations. *Ann R Coll Surg Engl* 1994; 76:95-98
23. Bennett-Guerrero E, Sorohan JG, Canada AT, Ayuso L, Newman MF, Reves JG, Mythen MG: e-aminocaproic acid plasma levels during cardiopulmonary bypass. *Anesth Analg* 1997; 85:248-251
24. Benoist S, Panis Y, Pannegeon V, Alves A, Valleur P: Predictive factors for perioperative blood transfusions in rectal resection for cancer: A multivariate analysis of a group of 212 patients. *Surgery* 2001; 129:433-439
25. Benoni G, Fredin H: Fibrinolytic inhibition with tranexamic acid reduces blood loss and blood transfusion after knee arthroplasty: a prospective, randomised, double-blind study of 86 patients. *J Bone Joint Surg Br* 1996; 78:434-440
26. Benoni G, Lethagen S, Nilsson P, Fredin H: Tranexamic acid, given at the end of the operation, does not reduce postoperative blood loss in hip arthroplasty. *Acta Orthop Scand* 2000; 71:250-254
27. Biesma DH, Marx JJ, Kraaijenhagen RJ, Franke W, Messinger D, van de Wiel A: Lower homologous blood requirement in autologous blood donors after treatment with recombinant human erythropoietin. *Lancet* 1994; 344:367-370
28. Billote DB, Glisson SN, Green D, Wixson RL: Efficacy of preoperative autologous blood donation: analysis of blood loss and transfusion practice in total hip replacement. *J Clin Anesth* 2000; 12:537-542
29. Bjessmo S, Ivert T: Blood loss after coronary artery bypass surgery: relations to patient variables and antithrombotic treatment. *Scand Cardiovasc J* 2000; 34:438-445
30. Blais RE, Hadjipavlou AG, Shulman G: Efficacy of autotransfusion in spine surgery: comparison of autotransfusion alone and with hemodilution and apheresis. *Spine* 1996; 21:2795-2800
31. Blauhut B, Harringer W, Bettelheim P, Doran JE, Spath P, Lundsgaard-Hansen P: Comparison of the effects of aprotinin and tranexamic acid on blood loss and related variables after cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1994; 108:1083-1091
32. Body SC, Birmingham J, Parks R, Ley C, Maddi R, Shernan SK, Siegel LC, Stover EP, D'Ambra MN, Levin J, Mangano DT, Spiess BD: Safety and efficacy of shed mediastinal blood transfusion after cardiac surgery: a multicenter observational study. Multicenter Study of Perioperative Ischemia Research Group. *J Cardiothorac Vasc Anesth* 1999; 13:410-416

33. Boldt J, Weber A, Mailer K, Papsdorf M, Schuster P: Acute normovolaemic haemodilution vs controlled hypotension for reducing the use of allogeneic blood in patients undergoing radical prostatectomy. *Br J Anaesth* 1999; 82:170-174
34. Bottner F, Pavone V, Johnson T, Heitkemper S, Sculco TP: Blood management after bilateral total knee arthroplasty. *Clin Orthop* 2003; 410:254-261
35. Boyer-Neumann C, Dreyfus M, Wolf M, Veyradier A, Meyer D: Multi-therapeutic approach to manage delivery in an alloimmunized patient with type 3 von Willebrand disease. *J Thromb Haemost* 2003; 1:190-192
36. Boylan JF, Klinck JR, Sandler AN, Arellano R, Greig PD, Nierenberg H, Roger SL, Glynn MF: Tranexamic acid reduces blood loss, transfusion requirements, and coagulation factor use in primary orthotopic liver transplantation. *Anesthesiology* 1996; 85:1043-1048
37. Braga M, Gianotti L, Vignali A, Gentilini O, Servida P, Bordignon C, Di Carlo V: Evaluation of recombinant human erythropoietin to facilitate autologous blood donation before surgery in anaemic patients with cancer of the gastrointestinal tract. *Br J Surg* 1995; 82:1637-1640
38. Brener SJ, Bhatt DL, Moliterno DJ, Schneider JP, Ellis SG, Topol EJ: Revisiting optimal anticoagulation with unfractionated heparin during coronary stent implantation. *Am J Cardiol* 2003; 92:1468-1471
39. Brown RS, Thwaites BK, Mongan PD: Tranexamic acid is effective in decreasing postoperative bleeding and transfusions in primary coronary artery bypass operations: a double-blind, randomized, placebo-controlled trial. *Anesth Analg* 1997; 85:963-970
40. Burman JF, Westlake AS, Davidson SJ, Rutherford LC, Rayner AS, Wright AM, Morgan CJ, Pepper JR: Study of five cell salvage machines in coronary artery surgery. *Transfus Med* 2002; 12:173-179
41. Cacciarelli TV, Keeffe EB, Moore DH, Burns W, Chuljian P, Busque S, Concepcion W, So SK, Esquivel CO: Primary liver transplantation without transfusion of red blood cells. *Surgery* 1996; 120:698-704
42. Cammerer U, Dietrich W, Rampf T, Braun SL, Richter JA: The predictive value of modified computerized thromboelastography and platelet function analysis for postoperative blood loss in routine cardiac surgery. *Anesth Analg* 2003; 96:51-57
43. Canver CC, Kroncke GM, Nichols RD, Murray EL, Mentzer RM Jr: The effects of intraoperative autologous whole blood sequestration on the need for transfusion of allogeneic blood and blood products in coronary bypass operations. *J Cardiovasc Surg (Torino)* 1995; 36:423-438
44. Capdevila X, Calvet Y, Biboulet P, Biron C, Rubenovitch J, d'Athis F: Aprotinin decreases blood loss and homologous transfusions in patients undergoing major orthopedic surgery. *Anesthesiology* 1998; 88:50-57
45. Casas JJ, Zuazu-Jausoro I, Mateo J, Oliver A, Litvan H, Muniz-Diaz E, Aris A, Caralps JM, Fontcuberta J: Aprotinin versus desmopressin for patients undergoing operations with cardiopulmonary bypass. A double-blind placebo-controlled study. *J Thorac Cardiovasc Surg* 1995; 110:1107-1117
46. Casati V, Bellotti F, Gerli C, Franco A, Oppizzi M, Cossolini M, Calori G, Benussi S, Alfieri O, Torri G: Tranexamic acid administration after cardiac surgery: A prospective, randomized, double-blind, placebo-controlled study. *Anesthesiology* 2001; 94:8-14
47. Casati V, Gerli C, Franco A, Torri G, D'Angelo A, Benussi S, Alfieri O: Tranexamic acid in off-pump coronary surgery: A preliminary, randomized, double-blind, placebo-controlled study. *Ann Thorac Surg* 2001; 72:470-475
48. Casati V, Speziali G, D'Alessandro C, Cianchi C, Antonietta Grasso M, Spagnolo S, Sandrelli L: Intraoperative low-volume acute normovolemic hemodilution in adult open-heart surgery. *Anesthesiology* 2002; 97:367-373

49. Cash KL, Brown T, Sausais L, Uehlinger J, Reed LJ: Severe delayed hemolytic transfusion reaction secondary to anti-At(a). *Transfusion* 1999; 39:834-837
50. Chan ACW, Blumgart LH, Wuest DL, Melendez JA, Fong Y: Use of preoperative autologous blood donation in liver resections for colorectal metastases. *Am J Surg* 1998; 175:461-465
51. Chanda A, Smith DR, Nanda A: Autotransfusion by cell saver technique in surgery of lumbar and thoracic spinal fusion with instrumentation. *J Neurosurg* 2002; 96(3 Suppl):298-303
52. Cheema P, El-Mefty O, Jazieh AR: Intraoperative haemorrhage associated with the use of extract of Saw Palmetto herb: a case report and review of the literature. *J Intern Med* 2001; 250:167-169
53. Chen L, Bracey AW, Radobancevic R, Cooper JR, Collard CD, Vaughn WK, Nussmeier NA: Clopidogrel and bleeding in patients undergoing elective coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 2004, 128:425-431
54. Chikada M, Furuse A, Kotsuka Y, Yagyu K: Open-heart surgery in Jehovah's Witness patients. *Cardiovasc Surg* 1996; 4:311-314
55. Cicek S, Demirkilic U, Kuralay E, Ozal E, Tatar H: Postoperative aprotinin: effect on blood loss and transfusion requirements in cardiac operations. *Ann Thorac Surg* 1996; 61:1372-1376
56. Cicek S, Demirkilic U, Ozal E, Kuralay E, Bingol H, Tatar H, Ozturk OY: Postoperative use of aprotinin in cardiac operations: an alternative to its prophylactic use. *J Thorac Cardiovasc Surg* 1996; 112:1462-1467
57. Cina CS, Bruin G: Acute normovolemic hemodilution (ANH) in surgery of the thoraco-abdominal aorta. A cohort study to evaluate coagulation parameters and blood products utilization. *J Cardiovasc Surg* 1999; 40:37-43
58. Cina CS, It SC, Clase CM, Bruin G: A cohort study of coagulation parameters and the use of blood products in surgery of the thoracic and thoracoabdominal aorta. *J Vasc Surg* 2001; 33:462-468
59. Clagett GP, Valentine RJ, Jackson MR, Mathison C, Kakish HB, Bengtson TD: A randomized trial of intraoperative autotransfusion during aortic surgery. *J Vasc Surg* 1999; 29:22-30
60. Clagett GP, Valentine RJ, Myers SI, Chervu A, Heller J: Does desmopressin improve hemostasis and reduce blood loss from aortic surgery? A randomized, double-blind study. *J Vasc Surg* 1995; 22:223-229
61. Clugston PA, Fitzpatrick DG, Kester DA, Folet B, Germann E: Autologous blood use in reduction mammoplasty: is it justified? *Plast Reconstr Surg* 1995; 95:824-828
62. Codisoti M, Mankad PS: Significant merits of a fibrin sealant in the presence of coagulopathy following paediatric cardiac surgery: randomised controlled trial. *Eur J Cardiothorac Surg* 2002; 22:200-205
63. Coffey A, Pittman J, Halbrook H, Fehrenbacher J, Beckman D, Hormuth D: The use of tranexamic acid to reduce postoperative bleeding following cardiac surgery: a double-blind randomized trial. *Am Surg* 1995; 61:566-568
64. Connelly NR, Kiessling BM, Brull SJ: Does tranexamic acid decrease bleeding in patients undergoing cardiopulmonary bypass? *Yale J Biol Med* 1994; 67:265-268
65. Connor JP, Morris PC, Alagoz T, Anderson B, Bottles K, Buller RE: Intraoperative autologous blood collection and autotransfusion in the surgical management of early cancers of the uterine cervix. *Obstet Gynecol* 1995; 86:373-378
66. Consten ECJ, Henny CP, Eijssman L, Dongelmans DA, van Oers MHJ: The routine use of fresh frozen plasma in operations with cardiopulmonary bypass is not justified. *J Thorac Cardiovasc Surg* 1996; 112:162-167
67. Copley LA, Richards BS, Safavi FZ, Newton PO: Hemodilution as a method to reduce transfusion requirements in adolescent spine fusion surgery. *Spine* 1999; 24:219-222

68. Daily PO, Lamphere JA, Dembitsky WP, Adamson RM, Dans NF: Effect of prophylactic epsilon-aminocaproic acid on blood loss and transfusion requirements in patients undergoing first-time coronary artery bypass grafting. A randomized, prospective, double-blind study. *J Thorac Cardiovasc Surg* 1994; 108:99-106
69. Dalmau A, Sabate A, Acosta F, Garcia-Huete L, Koo M, Reche M, Rafecas A, Figueras J, Jaurrieta E: Comparative study of antifibrinolytic drugs in orthotopic liver transplantation. *Transplant Proc* 1999; 31:2361-2362
70. Dalmau A, Sabate A, Acosta F, Garcia-Huete L, Koo M, Sansano T, Rafecas A, Figueras J, Jaurrieta E, Parrilla P: Tranexamic acid reduces red cell transfusion better than epsilon-aminocaproic acid or placebo in liver transplantation. *Anesth Analg* 2000; 91:29-34
71. Dalrymple-Hay MJ, Pack L, Deakin CD, Shephard S, Ohri SK, Haw MP, Livesey SA, Monro JL: Autotransfusion of washed shed mediastinal fluid decreases the requirement for autologous blood transfusion following cardiac surgery: a prospective randomized trial. *Eur J Cardiothorac Surg* 1999; 15:830-834
72. D'Ambra MN, Akins CW, Blackstone EH, Bonney SL, Cohn LH, Cosgrove DM, Levy JH, Lynch KE, Maddi R: Aprotinin in primary valve replacement and reconstruction: a multicenter, double-blind, placebo-controlled trial. *J Thorac Cardiovasc Surg* 1996; 112:1081-1089
73. Dave SP, Greenstein AJ, Sachar DB, Harpaz N, Aledort L: Bleeding diathesis in amyloidosis with renal insufficiency associated with Crohn's disease: response to desmopressin. *Am J Gastroenterol* 2002; 97:187-189
74. de Andrade JR, Jove M, Landon G, Frei D, Guilfoyle M, Young DC: Baseline hemoglobin as a predictor of risk of transfusion and response to Epoetin alfa in orthopedic surgery patients. *Am J Orthop* 1996; 25:533-542
75. de Pree C, Mermillod B, Hoffmeyer P, Beris P: Recombinant human erythropoietin as adjuvant treatment for autologous blood donation in elective surgery with large blood needs (≥ 5 units): a randomized study. *Transfusion* 1997; 37:708-714
76. del Pozo Pozo AI, Jimenez-Yuste V, Villar A, Quintana M, Hernandez-Navarro F: Successful thyroidectomy in a patient with Hermansky-Pudlak syndrome treated with recombinant activated factor VII and platelet concentrates. *Blood Coagul Fibrinolysis* 2002; 13:551-553
77. Despotis GJ, Levine V, Filos KS, Santoro SA, Joist JH, Spitznagel E, Goodnough LT: Evaluation of a new point-of-care test that measures PAF-mediated acceleration of coagulation in cardiac surgical patients. *Anesthesiology* 1996; 85:1311-1323
78. Despotis GJ, Levine V, Saleem R, Spitznagel E, Joist JH: Use of point-of-care test in identification of patients who can benefit from desmopressin during cardiac surgery: a randomised controlled trial. *Lancet* 1999; 354:106-110
79. Dich-Nielsen JO, Rajan RM, Jensen JJ: An anaphylactoid-like reaction following infusion of salvaged unwashed drain blood. *Can J Anaesth* 1998; 45:189
80. Dietrich W, Dilthey G, Spannagl M, Jochum M, Braun SL, Richter JA: Influence of high-dose aprotinin on anticoagulation, heparin requirement, and celite- and kaolin-activated clotting time in heparin-pretreated patients undergoing open-heart surgery. A double-blind, placebo-controlled study. *Anesthesiology* 1995; 83:679-689
81. Dignan RJ, Law DW, Seah PW, Manganas CW, Newman DC, Grant PW, Wolfenden HD: Ultra-low dose aprotinin decreases transfusion requirements and is cost effective in coronary operations. *Ann Thorac Surg* 2001; 71:158-63
82. Dryden PJ, O'Connor JP, Jamieson WR, Reid I, Ansley D, Sadeghi H, Burr LH, Munro AI, Merrick PM: Tranexamic acid reduces blood loss and transfusion in reoperative cardiac surgery. *Can J Anaesth* 1997; 44:934-941

83. Ehrlich M, Grabenwoger M, Cartes-Zumelzu F, Luckner D, Kovarik J, Laufer G, Kocher A, Konetschny R, Wolner E, Havel M: Operations on the thoracic aorta and hypothermic circulatory arrest: is aprotinin safe? *J Thorac Cardiovasc Surg* 1998; 115:220-225
84. Ekback G, Axelsson K, Rytberg L, Edlund B, Kjellberg J, Weckstrom J, Carlsson O, Schott U: Tranexamic acid reduces blood loss in total hip replacement surgery. *Anesth Analg* 2000; 91:1124-1130
85. Ekback G, Schott U, Axelsson K, Carlberg M: Perioperative autotransfusion and functional coagulation analysis in total hip replacement. *Acta Anaesthes Scand* 1995; 39:390-395
86. Engel JM, Hohaus T, Ruwoldt R, Menges T, Jurgensen I, Hempelmann G: Regional hemostatic status and blood requirements after total knee arthroplasty with and without tranexamic acid or aprotinin. *Anesth Analg* 2001; 92:775-780
87. Englberger L, Markart P, Eckstein FS, Immer FF, Berdat PA, Carrel TP: Aprotinin reduces blood loss in off-pump coronary artery bypass (OPCAB) surgery. *Eur J Cardiothorac Surg* 2002; 22:545-551
88. Engstrom KG, Appelblad M, Brorsson, B: Mechanisms behind operating room blood transfusions in coronary artery bypass graft surgery patients with insignificant bleeding. *J Cardiothorac Vasc Anes* 2002; 16:539-544
89. Ereth MH, Nuttall GA, Klindworth JT, MacVeigh I, Santrach PJ, Orszulak TA, Harmsen WS, Oliver W Jr: Does the platelet-activated clotting test (Hemo-STATUS) predict blood loss and platelet dysfunction associated with cardiopulmonary bypass? *Anesth Analg* 1997; 85:257-264
90. Ereth MH, Nuttall GA, Santrach PJ, Klindworth JT, Oliver WC Jr, Schaff HV: The relation between the platelet-activated clotting test (HemoSTATUS) and blood loss after cardiopulmonary bypass. *Anesthesiology* 1998; 88:962-969
91. Estella NM, Berry DL, Baker BW, Wali AT, Belfort MA: Normovolemic hemodilution before cesarean hysterectomy for placenta percreta. *Obstet Gynecol* 1997; 90:669-670
92. Etchason J, Petz L, Keeler E, Calhoun L, Kleinman S, Snider C, Fink A, Brook R: The cost effectiveness of preoperative autologous blood donations. *N Engl J Med* 1995; 332:719-724
93. Faris PM, Ritter MA, Abels RI: The effects of recombinant human erythropoietin on perioperative transfusion requirements in patients having a major orthopaedic operation. The American Erythropoietin Study Group. *J Bone Joint Surg Am* 1996; 78:62-72
94. Farrer A, Spark JI, Scott DJ: Autologous blood transfusion: the benefits to the patients undergoing abdominal aortic aneurysm repair. *J Vasc Nurs* 1997; 15:111-115
95. Feagan BG, Wong CJ, Kirkley A, Johnston DW, Smith FC, Whitsitt P, Wheeler SL, Lau CY: Erythropoietin with iron supplementation to prevent allogeneic blood transfusion in total hip joint arthroplasty. A randomized, controlled trial. *Ann Intern Med* 2000; 133:845-854
96. Ferraris VA, Ferraris SP, Joseph O, Wehner P, Mentzer RM Jr: Aspirin and postoperative bleeding after coronary artery bypass grafting. *Ann Surg* 2002; 235:820-827
97. Ferraz-Neto BH, Silva ED, Afonso RC, Gregory FH, Meira-Filho SP, Macedo CP, Parra OM, Leitao RM, Saad WA: Acute normovolemic intraoperative hemodilution in liver transplantation. *Transplant Proc* 1999; 31:3069-3070
98. Findlay JY, Rettke SR, Ereth MH, Plevak DJ, Krom RA, Kufner RP: Aprotinin reduces red blood cell transfusion in orthotopic liver transplantation: a prospective, randomized, double-blind study. *Liver Transpl* 2001; 7:802-807
99. Fitzsimmons MG, Peterfreund RA, Raines DE: Aprotinin administration and pulmonary thromboembolism during orthotopic liver transplantation: report of two cases. *Anesth Analg* 2001; 92:1418-1421
100. Florentino-Pineda I, Blakemore LC, Thompson GH, Poe-Kochert C, Adler P, Tripi P: The Effect of e-aminocaproic acid on perioperative blood loss in patients with idiopathic

- scoliosis undergoing posterior spinal fusion: a preliminary prospective study. *Spine* 2001; 26:1147-1151
101. Ford SM, Unsworth-White MJ, Aziz T, Tooze JA, van Besouw JP, Bevan DH, Treasure T: Platelet pheresis is not a useful adjunct to blood-sparing strategies in cardiac surgery. *J Cardiothorac Vasc Anesth* 2002; 16:321-329
 102. Fox JS, Amaranath L, Hoeltge GA, Andrish JT: Autologous blood transfusion and intraoperative cell salvage in a patient with homozygous sickle cell disease. *Cleve Clin J Med* 1994; 61:137-140
 103. Franchini M, de Gironcoli M, Lippi G, Manzato F, Aprili G, Gandini G: Prophylactic use of desmopressin in surgery of six patients with symptomatic heterozygous factor XI deficiency. *Haematologica* 2000; 85:106-107
 104. Friederich PW, Henny CP, Messelink EJ, Geerdink MG, Keller T, Kurth KH, Buller HR, Levi M: Effect of recombinant activated factor VII on perioperative blood loss in patients undergoing retropubic prostatectomy: a double-blind placebo-controlled randomised trial. *Lancet*. 2003; 361:201-205
 105. Garcia L, Sabate A, Domenech P, Martinez Brotons F, Drudis R, Jaurrieta E: Aprotinin in orthotopic liver transplantation. *Transplant Proc* 1995; 27:2290-2291
 106. Garcia-Huete L, Domenech P, Sabate A, Martinez-Brotons F, Jaurrieta E, Figueras J: The prophylactic effect of aprotinin on intraoperative bleeding in liver transplantation: a randomized clinical study. *Hepatology* 1997; 26:1143-1148
 107. Gardner A, Gibbs N, Evans C, Bell R: Relative cost of autologous red cell salvage versus allogeneic red cell transfusion during abdominal aortic aneurysm repair. *Anaesth Intensive Care* 2000; 28:646-649
 108. Gaudino M, Luciani N, Piancone FL, Bruno P, Rossi M, Schiavello R, Possati G: Perioperative management of a patient with Werlhof disease undergoing myocardial revascularization. *J Cardiovasc Surg (Torino)* 1999; 40:227-228
 109. Gelb AB, Roth RI, Levin J, London MJ, Noall RA, Hauck WW, Cloutier M, Verrier E, Mangano DT: Changes in blood coagulation during and following cardiopulmonary bypass. *Am J Clin Pathol* 1996; 106:87-99
 110. Gerrah R, Shargal Y, Elami A: Impaired oxygenation and increased hemolysis after cardiopulmonary bypass in patients with glucose-6-phosphate dehydrogenase deficiency. *Ann Thorac Surg* 2003; 76:523-527
 111. Gilbert JB, Malkowicz SB, Wein AJ: Cell saver and radical retropubic prostatectomy: analysis of cost-effectiveness. *Urology* 1995; 46:542-544
 112. Giurati G, Zanetta G, Caspani L, Landon F, Pellegrino A, Baglioni S, Mangioni C: Autologous blood transfusion in patients undergoing radical hysterectomy. *Br J Obstet Gynaecol* 1995; 102:64-65
 113. Glazier DB, Ciocca RG, Gosin JS, Murphy DP, Graham AM: Elective aortic surgery with minimal banked blood. *Am Surg* 1998; 64:171-174
 114. Godet G, Fleron MH, Vicaut E, Zubicki A, Bertrand M, Riou B, Kieffer E, Coriat P: Risk factors for acute postoperative renal failure in thoracic or thoracoabdominal aortic surgery: a prospective study. *Anesth Analg* 1997; 85:1227-1232
 115. Goh M, Klier CG, Kielczewski P, Wojno KJ, Kim K, Oesterling JE: Autologous blood donation prior to anatomical radical retropubic prostatectomy: is it necessary? *Urology* 1997; 49:569-573
 116. Goldstein DJ, DeRosa CM, Mongero LB, Weinberg AD, Michler RE, Rose EA, Oz MC, Smith CR: Safety and efficacy of aprotinin under conditions of deep hypothermia and circulatory arrest. *J Thorac Cardiovasc Surg* 1995; 110:1615-1621

117. Goldstein DJ, Oz MC, Smith CR, Friedlander JP, De Rosa CM, Mongero LB, Delphin E: Safety of repeat aprotinin administration for LVAD recipients undergoing cardiac transplantation. *Ann Thorac Surg* 1996; 61:692-695
118. Goldstein DJ, Seldomridge JA, Chen JM, Catanese KA, DeRosa CM, Weinberg AD, Smith CR, Rose EA, Levin HR, Oz MC: Use of aprotinin in LVAD recipients reduces blood loss, blood use, and perioperative mortality. *Ann Thorac Surg* 1995; 59:1063-1067
119. Gombotz H, Gries M, Sipurzynski S, Fruhwald S, Rehak P: Preoperative treatment with recombinant human erythropoietin or predeposit of autologous blood in women undergoing primary hip replacement. *Acta Anaesthesiol Scand* 2000; 44:737-742
120. Good L, Peterson E, Lisander B: Tranexamic acid decreases external blood loss but not hidden blood loss in total knee replacement. *Br J Anaesth* 2003; 90:596-599
121. Goodnough LT, Monk TG, Sicard G, Satterfield SA, Allen B, Anderson CB, Thompson RW, Flye W, Martin K: Intraoperative salvage in patients undergoing elective abdominal aortic aneurysm repair: an analysis of cost and benefit. *J Vasc Surg* 1996; 24:213-218
122. Goodnough LT, Price TH, Friedman KD, Johnston M, Ciavarella D, Khan N, Sacher R, Vogler WR, Wissel M, Abels RI: A phase III trial of recombinant human erythropoietin therapy in non-anemic orthopedic patients subjected to aggressive removal of blood for autologous use: Dose, response, toxicity, and efficacy. *Transfusion* 1994; 34:66-71
123. Gravlee GP, Arora S, Lavendar SW, Mills SA, Hudspeth AS, Cordell AR, James RL, Brockschmidt JK, Stuart JJ: Predictive value of blood clotting tests in cardiac surgical patients. *Ann Thorac Surg* 1994; 58:216-221
124. Gray CL, Amling CL, Polston GR, Powell CR, Kane CJ: Intraoperative cell salvage in radical retropubic prostatectomy. *Urology* 2001; 58:740-745
125. Grosvenor D, Goyal V, Goodman S: Efficacy of postoperative blood salvage following total hip arthroplasty in patients with and without deposited autologous units. *J Bone Joint Surg Am* 2000; 82-A:951-954
126. Hallak M, Sharon AS, Diukman R, Auslender R, Abramovici H: Supplementing iron intravenously in pregnancy. A way to avoid blood transfusions. *J Reprod Med* 1997; 42:99-103
127. Hardy JF, Belisle S, Dupont C, Harel F, Robitaille D, Roy M, Gagnon L: Prophylactic tranexamic acid and epsilon-aminocaproic acid for primary myocardial revascularization. *Ann Thorac Surg* 1998; 65:371-376
128. Harley BJ, Beaupre LA, Jones CA, Cinats JG, Guenther CR: The effect of epsilon aminocaproic acid on blood loss in patients who undergo primary total hip replacement: a pilot study. *Can J Surg* 2002; 45:185-190
129. Havel M, Grabenwoger F, Schneider J, Laufer G, Wollenek G, Owen A, Simon P, Teufelsbauer H, Wolner E: Aprotinin does not decrease early graft patency after coronary artery bypass grafting despite reducing postoperative bleeding and use of donated blood. *J Thorac Cardiovasc Surg* 1994; 107:807-810
130. Hayashi J, Kumon K, Takanashi S, Kawashima Y, Eguchi S, Takaku F, Yamamura H: Subcutaneous administration of recombinant human erythropoietin before cardiac surgery: a double-blind, multicenter trial in Japan. *Transfusion* 1994; 34:142-146
131. Hayes A, Murphy DB, McCarroll M: The efficacy of single-dose aprotinin 2 million KIU in reducing blood loss and its impact on the incidence of deep venous thrombosis in patients undergoing total hip replacement surgery. *J Clin Anesth* 1996; 8:357-360
132. Haynes SL, Torella F, Wong JC, Dalrymple K, James M, McCollum CN: Economic evaluation of a randomized clinical trial of haemodilution with cell salvage in aortic surgery. *Br J Surg* 2002; 89:731-736

133. Healy WL, Pfeifer BA, Kurtz SR, Johnson C, Johnson W, Johnston R, Sanders D, Karpman R, Hallack GN, Valeri CR: Evaluation of autologous shed blood for autotransfusion after orthopaedic surgery. *Clin Orthop* 1994; 299:53-59
134. Hedstrom M, Flordal PA, Ahl T, Svensson J, Dalen N: Autologous blood in hip replacement. No effect on blood loss but less increase of plasminogen activator inhibitor in a randomized series of 80 patients. *Acta Orthop Scand* 1996; 67:317-320
135. Heiss MM, Mempel W, Delanoff C, Jauch KW, Gabka C, Mempel M, Dieterich HJ, Eissner HJ, Schildberg FW: Blood transfusion-modulated tumor recurrence: first results of a randomized study of autologous versus allogeneic blood transfusion in colorectal cancer surgery. *J Clin Oncol* 1994; 12:1859-1867
136. Heiss MM, Tarabichi A, Delanoff C, Allgayer H, Jauch KW, Hernandez-Richter T, Mempel W, Beck KG, Schildberg FW, Messmer K: Perisurgical erythropoietin application in anemic patients with colorectal cancer: A double-blind randomized study. *Surgery* 1996; 119:523-527
137. Helm RE, Klemperer JD, Rosengart TK, Gold JP, Peterson P, DeBois W, Altorki NK, Lang S, Thomas S, Isom OW, Krieger KH: Intraoperative autologous blood donation preserves red cell mass but does not decrease postoperative bleeding. *Ann Thorac Surg* 1996; 62:1431-1441
138. Hendrice C, Schmartz D, Pradier O, Coron T, Kucharcezowski C, d'Hollander A, Capel P: Effects of aprotinin on blood loss, heparin monitoring tests, and heparin doses in patients undergoing coronary artery bypass surgery. *J Cardiothorac Vasc Anesth* 1995; 9:245-249
139. Hendriks HG, Meijer K, de Wolf JT, Klomp maker IJ, Porte RJ, de Kam PJ, Hagens AJ, Melsen T, Slooff MJ, van der Meer J: Reduced transfusion requirements by recombinant factor VIIa in orthotopic liver transplantation: a pilot study. *Transplantation* 2001;71:402-405
140. Hendriks HG, van der Meer J, Klomp maker IJ, Choudhury N, Hagens JA, Porte RJ, de Kam PJ, Slooff MJ, de Wolf JT: Blood loss in orthotopic liver transplantation: a retrospective analysis of transfusion requirements and the effects of autotransfusion of cell saver blood in 164 consecutive patients. *Blood Coagul Fibrinolysis* 2000; 11:S87-S93
141. Herregods L, Moerman A, Foubert L, Den Blauwen N, Mortier E, Poelaert J, Struys M: Limited intentional normovolemic hemodilution: ST-segment changes and use of homologous blood products in patients with left main coronary artery stenosis. *J Cardiothorac Vasc Anesth* 1997; 11:18-23
142. Hiippala S, Strid L, Wennerstrand M, Arvela V, Mantyla S, Ylinen J, Niemela H: Tranexamic acid (Cyklokapron) reduces perioperative blood loss associated with total knee arthroplasty. *Br J Anaesth* 1995; 74:534-537
143. Hiippala ST, Strid LJ, Wennerstrand MI, Arvela JV, Niemela HM, Mantyla SK, Kuisma RP, Ylinen JE: Tranexamic acid radically decreases blood loss and transfusions associated with total knee arthroplasty. *Anesth Analg* 1997; 84:839-844
144. Hirsch J, Wendt T, Kuhly P, Schaffartzik W: Point-of-care testing: measurement of coagulation. *Anaesthesia* 2001; 56:760-763
145. Hobisch-Hagen P, Schobersberger W, Falkensammer J, Luz G, Innerhofer P, Frischhut B, Puschendorf B, Mair J: No release of cardiac troponin I during major orthopedic surgery after acute normovolemic hemodilution. *Acta Anaesthesiol Scand* 1998; 42:799-804
146. Hohn L, Schweizer A, Licker M, Morel DR: Absence of beneficial effect of acute normovolemic hemodilution combined with aprotinin on allogeneic blood transfusion requirements in cardiac surgery. *Anesthesiology* 2002; 96:276-282

147. Holmes JH 4th, Jones MF, Anderson RP, Knopes KD, Guyton SW, Hall RA: The use of micro-dose aprotinin with continuous infusion in coronary artery bypass surgery. *J Cardiovasc Surg (Torino)* 1999; 40:621-626
148. Horlocker TT, Nuttall GA, Dekutoski MB, Bryant SC: The accuracy of coagulation tests during spinal fusion and instrumentation. *Anesth Analg* 2001; 93:33-38
149. Huber TS, Carlton LC, Irwin PB, Flug RR, Harward TR, Flynn TC, Seeger JM: Intraoperative autologous transfusion during elective infrarenal aortic reconstruction. *J Surg Res* 1997; 67:14-20
150. Husted H, Blond L, Sonne-Holm S, Holm G, Jacobsen TW, Gebuhr P: Tranexamic acid reduces blood loss and blood transfusions in primary total hip arthroplasty: a prospective randomized double-blind study in 40 patients. *Acta Orthop Scand* 2003; 74:665-669
151. Hyllner M, Avall A, Swolin B, Bengtson JP, Bengtsson A: Autologous blood transfusion in radical hysterectomy with and without erythropoietin therapy. *Obstet Gynecol* 2002; 99:757-762
152. Ivert T, Intonti M, Stain-Malmgren R, Dumitrescu A, Blomback M: Effects of aprotinin during cardiopulmonary bypass in patients treated with acetylsalicylic acid. *Scand Cardiovasc J* 1998; 32:289-295
153. Jansen AJ, Andreica S, Claeys M, D'Haese J, Camu F, Jochmans K: Use of tranexamic acid for an effective blood conservation strategy after total knee arthroplasty. *Br J Anaesth* 1999; 83:596-601
154. Janssens M, Joris J, David J, Lemaire R, Lamy M: High dose aprotinin reduces blood loss in patients undergoing total hip replacement surgery. *Anesthesiology* 1994; 80:23-29
155. Jeserschek R, Clar H, Aigner C, Rehak P, Primus B, Windhager R: Reduction of blood loss using high-dose aprotinin in major orthopaedic surgery: a prospective, double-blind, randomised and placebo-controlled study. *J Bone Joint Surg Br* 2003; 85:174-177
156. Jewell AE, Akowuah EF, Suvarna SK, Braidley P, Hopkinson D, Cooper G: A prospective randomised comparison of cardiotomy suction and cell saver for recycling shed blood during cardiac surgery. *Eur J Cardiothorac Surg* 2003; 23:633-636
157. Johnson HD, Morgan MS, Utley JR, Leyand SA, Nguyen-Duy T, Crawley DM: Comparative analysis of recovery of cardiopulmonary bypass residual blood: cell saver vs hemoconcentrator. *J Extra Corpor Technol* 1994; 26:194-199
158. Johnson LB, Plotkin JS, Kuo PC: Reduced transfusion requirements during major hepatic resection with use of intraoperative isovolemic hemodilution. *Am J Surg* 1998; 176:608-611
159. Jordan D, Delphin E, Rose E: Prophylactic epsilon-aminocaproic acid (EACA) administration minimizes blood replacement therapy during cardiac surgery. *Anesth Analg* 1995; 80:827-829
160. Kahraman S, Altunkaya H, Celebioglu B, Kanbak M, Pasaoglu I, Erdem K: The effect of acute normovolemic hemodilution on homologous blood requirements and total estimated red blood cell volume lost. *Acta Anaesthesiol Scand* 1997; 41:614-617
161. Kajikawa M, Nonami T, Kurokawa T, Hashimoto S, Harada A, Nakao A, Takagi H: Autologous blood transfusion for hepatectomy in patients with cirrhosis and hepatocellular carcinoma: use of recombinant human erythropoietin. *Surgery* 1994; 115:727-734
162. Karck M, Haverich A: Heart transplantation under coumarin therapy: friend or foe? *Eur J Anaesthesiol* 1994; 11:475-479
163. Karnezis TA, Stulberg SD, Wixson RL, Reilly P: The hemostatic effects of desmopressin on patients who had total joint arthroplasty. A double-blind randomized trial. *J Bone Joint Surg Am* 1994; 76:1545-1550

164. Karski JM, Teasdale SJ, Norman P, Carroll J, VanKessel K, Wong P, Glynn MF: Prevention of bleeding after cardiopulmonary bypass with high-dose tranexamic acid. Double-blind, randomized clinical trial. *J Thorac Cardiovasc Surg* 1995; 110:835-842
165. Kaspar M, Ramsay MA, Nguyen AT, Cogswell M, Hurst G, Ramsay KJ: Continuous small-dose tranexamic acid reduces fibrinolysis but not transfusion requirements during orthotopic liver transplantation. *Anesth Analg* 1997; 85:281-285
166. Kasper SM, Elsner F, Hilgers D, Grond S, Rutt J: A retrospective study of the effects of small-dose aprotinin on blood loss and transfusion needs during total hip arthroplasty. *Eur J Anaesthesiol* 1998; 15:669-675
167. Kasper SM, Giesecke T, Limpers P, Sabatowski R, Mehlhorn U, Diefenbach C: Failure of autologous fresh frozen plasma to reduce blood loss and transfusion requirements in coronary artery bypass surgery. *Anesthesiology* 2001; 95:81-86
168. Katsaros D, Petricevic M, Snow NJ, Woodhall DD, Van Bergen R: Tranexamic acid reduces postbypass blood use: a double-blinded, prospective, randomized study of 210 patients. *Ann Thorac Surg* 1996; 61:1131-1135
169. Katz K, Tamary H, Lahav J, Soudry M, Cohen IJ: Increased operative bleeding during orthopaedic surgery in patients with type I Gaucher disease and bone involvement. *Bull Hosp Jt Dis* 1999; 58:188-190
170. Kaufmann CR, Dwyer KM, Crews JD, Dols SJ, Trask AL: Usefulness of thrombelastography in assessment of trauma patient coagulation. *J Trauma* 1997; 42:716-722
171. Kemper RR, Menitove JE, Hanto DW: Cost analysis of intraoperative blood salvage during orthotopic liver transplantation. *Liver Transpl Surg* 1997; 3:513-517
172. Kesten S, de Hoyas A, Chaparro C, Westney G, Winton T, Maurer JR: Aprotinin reduces blood loss in lung transplant recipients. *Ann Thorac Surg* 1995; 59:877-879
173. Kettelhack C, Hones C, Messinger D, Schlag PM: Randomized multicentre trial of the influence of recombinant human erythropoietin on intraoperative and postoperative transfusion need in anaemic patients undergoing right hemicolectomy for carcinoma. *Br J Surg* 1998; 85:63-67
174. Kettner SC, Panzer OP, Kozek SA, Seibt FA, Stoiser B, Kofler J, Locker GJ, Zimpfer M: Use of abciximab-modified thrombelastography in patients undergoing cardiac surgery. *Anesth Analg* 1999; 89:580-584
175. Kilgore ML, Pacifico AD: Shed mediastinal blood transfusion after cardiac operations: a cost-effectiveness analysis. *Ann Thorac Surg* 1998; 65:1248-1254
176. Kjaergard HK, Trumbull HR: Vivostat system autologous fibrin sealant: preliminary study in elective coronary bypass grafting. *Ann Thorac Surg* 1998; 66:482-486
177. Klodell CT, Richardson JD, Bergamini TM, Spain DA: Does cell-saver blood administration and free hemoglobin load cause renal dysfunction? *Am Surg* 2001; 67:44-47
178. Kluger R, Olive DJ, Stewart AB, Blyth CM: Epsilon-aminocaproic acid in coronary artery bypass graft surgery: preincision or postheparin? *Anesthesiology* 2003; 99:1263-1269
179. Kochamba GS, Pfeffer TA, Sintek CF, Khonsari S: Intraoperative autotransfusion reduces blood loss after cardiopulmonary bypass. *Ann Thorac Surg* 1996; 61:900-903
180. Kohli N, Mallipeddi PK, Neff JM, Sze EH, Roat TW: Routine hematocrit after elective gynecologic surgery. *Obstet Gynecol* 2000; 95:847-850
181. Kosmadakis N, Messaris E, Maris A, Katsaragakis S, Leandros E, Konstadoulakis MM, Androulakis G: Perioperative erythropoietin administration in patients with gastrointestinal tract cancer: prospective randomized double-blind study. *Ann Surg* 2003; 237:417-421
182. Kyriss T, Wurst H, Friedel G, Jaki R, Toomes H: Reduced blood loss by aprotinin in thoracic surgical operations associated with high risk of bleeding. A placebo-controlled, randomized phase IV study. *Eur J Cardiothorac Surg* 2001; 20:38-41

183. Lab M, Welz A, Kocks M, Mayer G, Schwandt M, Hannekum A.: Aprotinin in elective primary bypass surgery: graft patency and clinical efficacy. *Eur J Cardiothorac Surg* 1995; 9:206-210
184. Langdown AJ, Field J, Grote J, Himayat H: Aprotinin (Trasylol) does not reduce bleeding in primary total hip arthroplasty. *J Arthroplasty* 2000; 15:1009-1012
185. Laub GW, Riebman JB, Chen C, Adkins MS, Anderson WA, Fernandez J, McGrath LB: The impact of aprotinin on coronary artery bypass graft patency. *Chest* 1994; 106:1370-1375
186. Lazzara RR, Kidwell FE, Kraemer MF, Wood JA, Starr A: Reduction in costs, blood products, and operating time in patients undergoing open heart surgery. *Arch Surg* 1997; 132:858-860
187. Le Roux PD, Elliott JP, Winn HR: Blood transfusion during aneurysm surgery. *Neurosurgery* 2001; 49:1068-1075
188. Lemmer JH Jr, Dilling EW, Morton JR, Rich JB, Robicsek F, Bricker DL, Hantler CB, Copeland JG 3rd, Ochsner JL, Daily PO, Whitten CW, Noon GP, Maddi R: Aprotinin for primary coronary artery bypass grafting: a multicenter trial of three dose regimens. *Ann Thorac Surg* 1996; 62:1659-1667
189. Lemmer JH Jr, Metzdorff MT, Krause AH Jr, Martin MA, Okies JE, Hill JG: Emergency coronary artery bypass graft surgery in abciximab-treated patients. *Ann Thorac Surg* 2000; 69:90-95
190. Lemmer JH Jr, Stanford W, Bonney SL, Breen JF, Chomka EV, Eldredge WJ, Holt WW, Karp RB, Laub GW, Lipton MJ, Schaff HV, Tatoes CJ, Rumberger JA: Aprotinin for coronary bypass operations: efficacy, safety, and influence on early saphenous vein graft patency. A multicenter, randomized, double-blind, placebo-controlled study. *J Thorac Cardiovasc Surg* 1994; 107:543-551
191. Lentschener C, Benhamou D, Mercier FJ, Boyer-Neumann C, Naueau S, Smadja C, Wold M, Franco D: Aprotinin reduces blood loss and transfusion requirement in patients undergoing elective liver resection. *Anesth Analg* 1997; 84:875-881
192. Lentschener C, Cottin P, Bouaziz H, Mercier FJ, Wolf M, Aljabi Y, Boyer-Neumann C, Benhamou D: Reduction of blood loss and transfusion requirement by aprotinin in posterior lumbar spine fusion. *Anesth Analg* 1999; 89:590-597
193. Levy JH, Pifarre R, Schaff HV, Horrow JC, Albus R, Spiess B, Rosengart TK, Murray J, Clark RE, Smith P, Nadel A, Bonney SL, Kleinfeld R: A multicenter, double-blind, placebo-controlled trial of aprotinin for reducing blood loss and the requirement for donor-blood transfusion in patients undergoing repeat coronary artery bypass grafting. *Circulation* 1995; 92:2236-2244
194. Levy O, Martinowitz U, Oran A, Tauber C, Horoszowski H: The use of fibrin tissue adhesive to reduce blood loss and the need for blood transfusion after total knee arthroplasty. A prospective, randomized, multicenter study. *J Bone Joint Surg Am* 1999; 81:1580-1588
195. Lindgren L, Yli-Hankala A, Halme L, Koskimies S, Orko R: Transfusion-related acute lung injury (TRALI) after fresh frozen plasma in a patient with coagulopathy. *Acta Anaesthesiol Scand* 1996; 40:641-644
196. Lisander B, Jonsson R, Nordwall A: Combination of blood-saving methods decreases homologous blood requirements in scoliosis surgery. *Anaesth Intensive Care* 1996; 24:555-558
197. Llamas P, Cabrera R, Gomez-Arnau J, Fernandez MN: Hemostasis and blood requirements in orthotopic liver transplantation with and without high-dose aprotinin. *Haematologica* 1998; 83:338-346
198. Loubser PG: Use of acute normovolemic hemodilution for blood conservation during off-pump coronary artery bypass surgery. *Can J Anaesth* 2003; 50:620-621 Erratum in: *Can J Anaesth*. 2003 Aug-Sep;50(7):754

199. Mackay Z, Shugufta Q, Din M, Guru AA: Hemodilution in complicated high velocity vascular injuries of limbs. *J Cardiovasc Surg* 1996; 37:217-221
200. Makrydimas G, Lolis D, Lialios G, Tsiara S, Georgiou I, Bourantas KL: Recombinant human erythropoietin treatment of postpartum anemia. Preliminary results. *Eur J Obstet Gynecol Reprod Biol* 1998; 81:27-31
201. Marcel RJ, Stegall WC, Suit CT, Arnold JC, Vera RL, Ramsay MA, O'Donnell MB, Swygert TH, Hein AT, Whitten CW: Continuous small dose aprotinin controls fibrinolysis during orthotopic liver transplantation. *Anesth Analg* 1996; 82:1122-1125
202. Marty-Ane CH, Alric P, Picot MC, Picard E, Colson P, Mary H: Ruptured abdominal aortic aneurysm: influence of intraoperative management on surgical outcome. *J Vasc Surg* 1995; 22:780-786
203. Matot I, Scheinin O, Jurim O, Eid A: Effectiveness of acute normovolemic hemodilution to minimize allogeneic blood transfusion in major liver resections. *Anesthesiology* 2002; 97:794-800
204. McGill N, O'Shaughnessy D, Pickering R, Herbertson M, Gill R: Mechanical methods of reducing blood transfusion in cardiac surgery: randomised controlled trial. *BMJ* 2002; 324:1299-303
205. Menges T, Wagner RM, Welters I, Ruwoldt R, Boldt J, Hempelmann G: The role of protein C-thrombomodulin system and fibrinolysis during cardiovascular surgery: influence of acute preoperative plasmapheresis. *J Cardiothorac Vasc Anesth* 1996; 10:482-489
206. Menges T, Welters I, Wagner RM, Boldt J, Dapper F, Hempelmann G: The influence of acute preoperative plasmapheresis on coagulation tests, fibrinolysis, blood loss and transfusion requirements in cardiac surgery. *Eur J Cardiothorac Surg* 1997; 11:557-563
207. Menichetti A, Tritapepe L, Ruvolo G, Speziale G, Cogliati A, Di Giovanni C, Pacilli M, Criniti A: Changes in coagulation patterns, blood loss and blood use after cardiopulmonary bypass: aprotinin vs tranexamic acid vs epsilon aminocaproic acid. *J Cardiovasc Surg* 1996; 37:401-407
208. Mercuriali F, Gualtieri G, Sinigaglia L, Inghilleri G, Biffi E, Vinci A, Colotti MT, Barosi G, Lambertengh Delilieri G: Use of recombinant human erythropoietin to assist autologous blood donation by anemic rheumatoid arthritis patients undergoing major orthopedic surgery. *Transfusion* 1994; 34:501-506
209. Mercuriali F, Inghilleri G, Biffi E, Colotti MT, Vinci A, Oriani G: Epoetin alfa in low hematocrit patients to facilitate autologous blood donation in total hip replacement: a randomized, double-blind, placebo-controlled, dose-ranging study. *Acta Haematol* 1998; 100:69-76
210. Molenaar IQ, Legnani C, Groenland TH, Palareti G, Begliomini B, Terpstra OT, Porte RJ: Aprotinin in orthotopic liver transplantation: evidence for a prohemostatic, but not a prothrombotic, effect. *Liver Transpl* 2001; 7:896-903
211. Mongan PD, Brown RS, Thwaites BK: Tranexamic acid and aprotinin reduce postoperative bleeding and transfusions during primary coronary revascularization. *Anesth Analg* 1998; 87:258-265
212. Monte S, Lyons G: Peripartum management of a patient with Glanzmann's thrombasthenia using Thrombelastograph. *Br J Anaesth* 2002; 88:734-738
213. Mora Mangano CT, Neville MJ, Hsu PH, Mignea I, King J, Miller DC: Aprotinin, blood loss, and renal dysfunction in deep hypothermic circulatory arrest. *Circulation* 2001; 104:I276-1281
214. Moscardo F, Perez F, de la Rubia J, Balerdi B, Lorenzo JI, Senent ML, Aznar I, Carceller S, Sanz MA: Successful treatment of severe intra-abdominal bleeding associated with

- disseminated intravascular coagulation using recombinant activated factor VII. *Br J Haematol* 2001; 114:174-176
215. Muller U, Roder C, Pisan M, Orlor R, El-Kerdi A, Eggli S: Autologous blood donation in total knee arthroplasties is not necessary. *Acta Orthop Scand* 2004 ;75:66-70
 216. Murkin JM, Haig GM, Beer KJ, Cicutti N, McCutchen J, Comunale ME, Hall R, Ruzicka BB: Aprotinin decreases exposure to allogeneic blood during primary unilateral total hip replacement. *J Bone Joint Surg Am* 2000; 82:675-684
 217. Murkin JM, Lux J, Shannon NA, Guiraudon GM, Menkis AH, McKenzie FN, Novick RJ: Aprotinin significantly decreases bleeding and transfusion requirements in patients receiving aspirin and undergoing cardiac operations. *J Thorac Cardiovasc Surg* 1994; 107:554-561
 218. Murkin JM, Shannon NA, Bourne RB, Rorabeck CH, Cruickshank M, Wyile G: Aprotinin decreases blood loss in patients undergoing revision or bilateral total hip arthroplasty. *Anesth Analg* 1995; 80:343-348
 219. Murray DJ, Pennell BJ, Weinstein SL, Olson JD: Packed red cells in acute blood loss: Dilutional coagulopathy as a cause of surgical bleeding. *Anesth Analg* 1995; 80:336-342
 220. Myers ER, Clarke-Pearson DL, Olt GJ, Soper JT, Berchuck A: Preoperative coagulation testing on a gynecologic oncology service. *Obstet Gynecol* 1994; 83:438-444
 221. Nash J, Hodgson C, Desmond M, Russell GN, Jackson M: Routine use of a cell saver during coronary artery by-pass grafting. *Transfusion* 1996; 36:657-658
 222. Neilipovitz DT, Murto K, Hall L, Barrowman NJ, Splinter WM: A randomized trial of tranexamic acid to reduce blood transfusion for scoliosis surgery. *Anesth Analg* 2001; 93:82-87
 223. Nuttall GA, Oliver WC, Beynen FM, Santrach PJ, Strickland RA, Murray MJ: Determination of normal versus abnormal activated partial thromboplastin time and prothrombin time after cardiopulmonary bypass. *J Cardiothorac Vasc Anesth* 1995; 9:355-361
 224. Nuttall GA, Oliver WC, Ereth MH, Santrach PJ, Bryant SC, Orszulak TA, Schaff HV: Comparison of blood-conservation strategies in cardiac surgery patients at high risk for bleeding. *Anesthesiology* 2000; 92:674-682
 225. Nuttall GA, Oliver WC, Ereth MH, Santrach PJ: Coagulation tests predict bleeding after cardiopulmonary bypass. *J Cardiothorac Vasc Anesth* 1997; 11:815-823
 226. Oefelein MG, Colangelo LA, Rademaker AW, McVary KT: Intraoperative blood loss and prognosis in prostate cancer patients undergoing radical retropubic prostatectomy. *J Urol* 1995; 154:442-447
 227. O'Hara PJ, Hertzner NR, Krajewski LP, Cox GS, Beven EG: Reduction in the homologous blood requirement for abdominal aortic aneurysm repair by the use of preadmission autologous blood donation. *Surgery* 1994; 115:69-76
 228. Oliver WC Jr, Santrach PJ, Danielson GK, Nuttall GA, Schroeder DR, Ereth MH: Desmopressin does not reduce bleeding and transfusion requirements in congenital heart operations. *Ann Thorac Surg* 2000; 70:1923-1930
 229. Olsfanger D, Fredman B, Goldstein B, Shapiro A, Jedeikin R: Acute normovolaemic haemodilution decreases postoperative allogeneic blood transfusion after total knee replacement. *Br J Anaesth* 1997; 79:317-321
 230. Ozkisacik E, Islamoglu F, Posacioglu H, Yagdi T, Basarir S, Omay SB, Ozbaran M, Buket S: Desmopressin usage in elective cardiac surgery. *J Cardiovasc Surg (Torino)* 2001; 42:741-747
 231. Pagani FD, Polito RJ, Bolling SF: Mitral valve reconstruction in sickle cell disease. *Ann Thorac Surg* 1996; 61:1841-1843
 232. Palmer JD, Francis DA, Roath OS, Francis JL, Iannotti F: Hyperfibrinolysis during intracranial surgery: effect of high dose aprotinin. *J Neurosurg Psychiatry* 1995; 58:104-106

233. Palmer JD, Francis JL, Pickard JD, Iannotti F: The efficacy and safety of aprotinin for hemostasis during intracranial surgery. *J Neurosurg* 2003; 98:1208-1216
234. Parolari A, Antona C, Alamanni F, Spirito R, Naliato M, Gerometta P, Arena V, Biglioli P: Aprotinin and deep hypothermic circulatory arrest: there are no benefits even when appropriate amounts of heparin are given. *Eur J Cardiothorac Surg* 1997; 11:149-156
235. Patrassi GM, Viero M, Sartori MT, De Silvestro G, Rossaro L, Burra P, Nolli ML, Piccinni P, Bassi N: Aprotinin efficacy on intraoperative bleeding and transfusion requirements in orthotopic liver transplantation. *Transfusion* 1994; 34:507-511
236. Pellegrino A, Landoni F, Cormio G, Ferrando P, Lissoni A, Scalabrino S, Mangioni C: Effectiveness of autologous blood transfusion in patients undergoing radical hysterectomy. *Ann Chir Gynaecol* 1995; 84:391-394
237. Penta de Peppo A, Pierri MD, Scafuri A, De Paulis R, Colantuono G, Caprara E, Tomai F, Chiariello L: Intraoperative antifibrinolysis and blood-saving techniques in cardiac surgery. Prospective trial of 3 antifibrinolytic drugs. *Tex Heart Inst J* 1995; 22:231-236
238. Pinkerton PH, Covens A: Autologous blood transfusion in radical hysterectomy. *Transfus Med* 1996; 6:223-225
239. Pirat A, Sargin D, Torgay A, Arslan G: Identification of preoperative predictors of intraoperative blood transfusion requirement in orthotopic liver transplantation. *Transplant Proc* 2002; 34:2153-2155
240. Pleym H, Stenseth R, Wahba A, Bjella L, Karevold A, Dale O: Single-dose tranexamic acid reduces postoperative bleeding after coronary surgery in patients treated with aspirin until surgery. *Anesth Analg* 2003; 96:923-928
241. Pleym H, Stenseth R, Wahba A, Bjella L, Tromsdal A, Karevold A, Dale O: Prophylactic treatment with desmopressin does not reduce postoperative bleeding after coronary surgery in patients treated with aspirin before surgery. *Anesth Analg* 2004; 98:578-584
242. Porte RJ, Molenaar IQ, Begliomini B, Groenland TH, Januszkiewicz A, Lindgren L, Palareti G, Hermans J, Terpstra OT: Aprotinin and transfusion requirements in orthotopic liver transplantation: a multicentre randomised double-blind study. *Lancet* 2000; 355:1303-1309
243. Posacioglu H, Apaydin AZ, Islamoglu F, Calkavur T, Yagdi T, Atay Y, Buket S: Adverse effects of cell saver in patients undergoing ruptured abdominal aortic aneurysm repair. *Ann Vasc Surg* 2002; 16:450-455
244. Potter PS, Waters JH, Burger GA, Mraovic B: Application of cell-salvage during cesarean section. *Anesthesiology* 1999; 90:619-621
245. Price TH, Goodnough LT, Vogler W, Sacher RA, Hellman RM, Johnston MF, Bolgiano DC, Abels RI: The effect of recombinant human erythropoietin on the efficacy of autologous blood donation in patients with low hematocrits: A multicenter, randomized, double-blind controlled trial. *Transfusion* 1996; 36:29-36
246. Propst JW, Siegel LC, Feeley TW: Effect of aprotinin on transfusion requirements during repeat sternotomy for cardiac transplantation surgery. *Transplant Proc* 1994; 26:3719-3721
247. Pugh SC, Wielogorski AK: A comparison of the effects of tranexamic acid and low-dose aprotinin on blood loss and homologous blood usage in patients undergoing cardiac surgery. *J Cardiothorac Vasc Anesth* 1995; 9:240-244
248. Qvist N, Boesby S, Wolff B, Hansen CP: Recombinant human erythropoietin and hemoglobin concentration at operation and during the postoperative period: reduced need for blood transfusions in patients undergoing colorectal surgery--prospective double-blind placebo-controlled study. *World J Surg* 1999; 23:30-35
249. Rainaldi MP, Tazzari PL, Scagliarini G, Borghi B, Conte R: Blood salvage during caesarean section. *Br J Anaesth* 1998; 80:195-198

250. Ramnath AN, Naber HR, de Boer A, Leusink JA: No benefit of intraoperative whole blood sequestration and autotransfusion during coronary artery bypass grafting: results of a randomized clinical trial. *J Thorac Cardiovasc Surg* 2003; 125:1432-1437
251. Ranaboldo CJ, Thompson JF, Davies JN, Shutt AM, Francis JN, Roath OS, Webster JH, Chant AD: Prospective randomized placebo-controlled trial of aprotinin for elective aortic reconstruction. *Br J Surg* 1997; 84:1110-1113
252. Rauh MA, Bayers-Thering M, LaButti RS, Krackow KA: Preoperative administration of epoetin alfa to total joint arthroplasty patients. *Orthopedics* 2002; 25:317-320
253. Ray JG, Deniz S, Olivieri A, Pollex E, Vermeulen MJ, Alexander KS, Cain DJ, Cybulsky I, Hamielec CM: Increased blood product use among coronary artery bypass patients prescribed preoperative aspirin and clopidogrel. *BMC Cardiovasc Disord* 2003; 3:3
254. Ray MJ, Hales MM, Brown L, O'Brien MF, Stafford EG: Postoperatively administered aprotinin or epsilon aminocaproic acid after cardiopulmonary bypass has limited benefit. *Ann Thorac Surg* 2001; 72:521-526
255. Ray MJ, Marsh NA, Just SJ, Perrin EJ, O'Brien MF, Hawson GA: Preoperative platelet dysfunction increases the benefit of aprotinin in cardiopulmonary bypass. *Ann Thorac Surg* 1997; 63:57-63
256. Rebarber A, Lonser R, Jackson S, Copel JA, Sipes S: The safety of intraoperative autologous blood collection and autotransfusion during cesarean section. *Am J Obstet Gynecol* 1998; 179:715-720
257. Reents W, Babin-Ebell J, Misoph MR, Schwarzkopf A, Elert O: Influence of different autotransfusion devices on the quality of salvaged blood. *Ann Thorac Surg* 1999; 68:58-62
258. Rehm M, Orth V, Kreimeier U, Thiel M, Haller M, Brechtelsbauer H, Finsterer U: Changes in intravascular volume during acute normovolemic hemodilution and intraoperative retransfusion in patients with radical hysterectomy. *Anesthesiology* 2000; 92:657-664
259. Reich DL, Yanakakis MJ, Vela-Cantos FP, DePerio M, Jacobs E: Comparison of bedside coagulation monitoring tests with standard laboratory tests in patients after cardiac surgery. *Anesth Analg* 1999; 88:312-319
260. Riou B, Arock M, Guerrero M, Ramos M, Thoreux P, Guillosson JJ, Roy-Camille R, Viars P: Haematological effects of postoperative autotransfusion in spinal surgery. *Acta Anaesthesiol Scand* 1994; 38:336-341
261. Roberts WA, Kirkley SA, Newby M: A cost comparison of allogeneic and preoperatively or intraoperatively donated autologous blood. *Anesth Analg* 1996; 83:129-133
262. Robinson J, Nawaz S, Beard JD: Randomized, multicentre, double-blind, placebo-controlled trial of the use of aprotinin in the repair of ruptured abdominal aortic aneurysm. *Br J Surg* 2000; 87:754-757
263. Rocha E, Hidalgo F, Llorens R, Melero JM, Arroyo JL, Paramo JA: Randomized study of aprotinin and DDAVP to reduce postoperative bleeding after cardiopulmonary bypass surgery. *Circulation* 1994; 90:921-927
264. Rollo VJ, Hozack WJ, Rothman RH, Chao W, Eng KO: Prospective randomized evaluation of blood salvage techniques for primary total hip arthroplasty. *J Arthroplasty* 1995; 10:532-539
265. Rousou JA, Engelman RM, Flack JE 3rd, Deaton DW, Owen SG: Tranexamic acid significantly reduces blood loss associated with coronary revascularization. *Ann Thorac Surg* 1995; 59:671-675
266. Rzyman W, Dziadziuszko R, Skokowski J, Wilimski R, Raiter A, Szymanowska A, Jassem J: The influence of blood transfusion on survival in operated non-small cell lung cancer patients. *J Thorac Cardiovasc Surg* 2003; 126:755-760
267. Safwat AM, Bush R, Prevec W, Reitan JA: Intraoperative use of platelet-plasmapheresis in vascular surgery. *J Clin Anesth* 2002; 14:10-14

268. Samama CM, Langeron O, Rosencher N, Capdevila X, Rouche P, Pegoix M, Berniere J, Coriat P: Aprotinin versus placebo in major orthopedic surgery: a randomized, double-blinded, dose-ranging study. *Anesth Analg* 2002; 95:287-293
269. Samama CM, Quezada R, Riou B, Mourgeon E, Arock M, Ankri A, Viars P: Intraoperative measurement of activated partial thromboplastin time and prothrombin time with a new compact monitor. *Acta Anaesth Scand* 1994; 38:232-237
270. Sandrelli L, Pardini A, Lorusso R, Sala ML, Licenziati M, Alfieri O: Impact of autologous blood predonation on a comprehensive blood conservation program. *Ann Thorac Surg* 1995; 59:730-755
271. Santamaria A, Mateo J, Oliver A, Litvan H, Murillo J, Souto JC, Fontcuberta J: The effect of two different doses of aprotinin on hemostasis in cardiopulmonary bypass surgery: similar transfusion requirements and blood loss. *Haematologica* 2000; 85:1277-1284
272. Schmied H, Schiferer A, Sessler DI, Meznik C: The effects of red-cell scavenging, hemodilution, and active warming on allogenic blood requirements in patients undergoing hip or knee arthroplasty. *Anesth Analg* 1998; 86:387-391
273. Scott SN, Boeve TJ, McCulloch TM, Fitzpatrick KA, Karnell LH: The effects of epoetin alfa on transfusion requirements in head and neck cancer patients: a prospective, randomized, placebo-controlled study. *Laryngoscope* 2002; 112:1221-1229
274. Scudamore CH, Randall TE, Jewesson PJ, Shackleton CR, Salvian AJ, Fagan M, Frighetto L, Growe GH, Scarth I, Erb SR: Aprotinin reduces the need for blood products during liver transplantation. *Am J Surg* 1995; 169:546-549
275. Seigne PW, Shorten GD, Johnson RG, Comunale ME: The effects of aprotinin on blood product transfusion associated with thoracic aortic surgery requiring deep hypothermic circulatory arrest. *J Cardiothorac Vasc Anesth* 2000; 14:676-681
276. Serrano FJ, Monux G, Aroca M: Should the cell saver autotransfusion system be routinely used in elective aortic surgery? *Ann Vasc Surg* 2000; 14:663-668
277. Shapiro GS, Boachie-Adjei O, Dhawlikar SH, Maier LS: The use of Epoetin alfa in complex spine deformity surgery. *Spine* 2002; 27:2067-2071
278. Shenolikar A, Wareham K, Newington D, Thomas D, Hughes J, Downes M: Cell salvage autotransfusion in total knee replacement surgery. *Transfus Med* 1997; 7:277-280
279. Sheridan DP, Card RT, Pinilla JC, Harding SM, Thomson DJ, Gauthier L, Drotar D: Use of desmopressin acetate to reduce blood transfusion requirements during cardiac surgery in patients with acetylsalicylic-acid-induced platelet dysfunction. *Can J Surg* 1994; 37:33-36
280. Shigeta O, Kojima H, Jikuya T, Terada Y, Atsuni N, Sakakibara Y, Nagasawa T, Mitsui T: Aprotinin inhibits plasmin-induced platelet activation during cardiopulmonary bypass. *Circulation* 1997; 96:569-574
281. Shore-Lesserson L, Reich DL, DePerio M, Silvay G: Autologous platelet-rich plasmapheresis: risk versus benefit in repeat cardiac operations. *Anesth Analg* 1995; 81:229-235
282. Shore-Lesserson L, Reich DL, Vela-Cantos F, Ammar T, Ergin MA: Tranexamic acid reduces transfusions and mediastinal drainage in repeat cardiac surgery. *Anesth Analg* 1996; 83:18-26
283. Shulman G, Grecula MJ, Hadjipavlou AG: Intraoperative autotransfusion in hip arthroplasty. *Clin Orthop* 2002; 396:119-30
284. Shulman G, Solanki DR, Hadjipavlou A: Augmented autologous transfusions in major reconstructive spine surgery. *J Clin Apheresis* 1998; 13:62-68
285. Simonian PT, Robinson RP: Titanium contamination of recycled Cell Saver blood in revision hip arthroplasty. *J Arthroplasty* 1995; 10:83-86

286. Slappendel R, Huvers FC, Benraad B, Novakova I, van Hellenmond GG: Use of recombinant factor VIIa to reduce postoperative bleeding after total hip arthroplasty in a patient with cirrhosis and thrombocytopenia. *Anesthesiology* 2002; 96:1525-1527
287. Slaughter TF, Faghih F, Greenberg CS, Leslie JB, Sladen RN: The effects of epsilon-aminocaproic acid on fibrinolysis and thrombin generation during cardiac surgery. *Anesth Analg* 1997; 85:1221-1226
288. Sokolic V, Bukovic D, Fures R, Zadro M, Scuric I, Colak F, Celovec K, Bocan A: Recombinant factor VIIa (rFVIIa) is effective at massive bleeding after caesarean section--a case report. *Coll Antropol* 2002; 26:155-157
289. Sowade O, Warnke H, Scigalla P, Sowade B, Franke W, Messinger D, Gross J: Avoidance of allogeneic blood transfusions by treatment with epoetin beta (recombinant human erythropoietin) in patients undergoing open-heart surgery. *Blood* 1997; 89:411-418
290. Sowade O, Ziemer S, Sowade B, Franke W, Messinger D, Ziebell E, Scigalla P, Warnke H: The effect of preoperative recombinant human erythropoietin therapy on platelets and hemostasis in patients undergoing cardiac surgery. *J Lab Clin Med* 1997; 129:376-383
291. Spark JI, Chetter IC, Kester RC, Scott DJ: Allogeneic versus autologous blood during abdominal aortic aneurysm surgery. *Eur J Vasc Endovasc Surg* 1997; 14:482-486
292. Speekenbrink RG, Wildevuur CR, Sturk A, Eijnsman L: Low-dose and high-dose aprotinin improve hemostasis in coronary operations. *J Thorac Cardiovasc Surg* 1996; 112:523-530
293. Spiess BD, Gillies BSA, Chandler W, Verrier E: Changes in transfusion therapy and reexploration rate after institution of a blood management program in cardiac surgical patients. *J Cardiothorac Vasc Anesth* 1995; 9:168-173
294. Stewart A, Newman L, Sneddon K, Harris M: Aprotinin reduces blood loss and the need for transfusion in orthognathic surgery. *Br J Oral Maxillofac Surg* 2001; 39:365-370
295. Stover EP, Siegel LC, Hood PA, O'Riordan GE, McKenna TR: Platelet-rich plasma sequestration, with therapeutic platelet yields, reduces allogeneic transfusion in complex cardiac surgery. *Anesth Analg* 2000; 90:509-516
296. Stratmann G, Russell IA, Merrick SH: Use of recombinant factor VIIa as a rescue treatment for intractable bleeding following repeat aortic arch repair. *Ann Thorac Surg* 2003; 76:2094-2097
297. Sutton SW, Hunley EK, Duncan MA, Rodriguez R, Meyers TP: Sickle cell disease and aortic valve replacement: use of cardiopulmonary bypass, partial exchange transfusion, platelet sequestration, and continuous hemofiltration. *Tex Heart Inst J* 1999; 26:283-288
298. Swart MJ, Gordon PC, Hayse-Gregson PB, Dyer RA, Swanepoel AL, Buckels NJ, Schall R, Odell JA: High-dose aprotinin in cardiac surgery--a prospective, randomized study. *Anaesth Intensive Care* 1994; 22:529-533
299. Szalay D, Wong D, Lindsay T: Impact of red cell salvage on transfusion requirements during elective abdominal aortic aneurysm repair. *Ann Vasc Surg* 1999; 13:576-581
300. Tabuchi N, Njo TL, Tigchelaar I, Huyzen RJ, Boonstra PW, van Oeveren W: Monitoring of anticoagulation in aprotinin-treated patients during heart operation. *Ann Thorac surg* 1994; 58:774-777
301. Taggart DP, Djapardy V, Naik M, Davies A: A randomized trial of aprotinin (Trasylol) on blood loss, blood product requirement, and myocardial injury in total arterial grafting. *J Thorac Cardiovasc Surg* 2003; 126:1087-1094
302. Tawes RL Jr, Duvall TB: Is the "salvaged-cell syndrome" myth or reality? *Am J Surg* 1996; 172:172-174
303. Temeck BK, Bachenheimer LC, Katz NM, Coughlin SS, Wallace RB: Desmopressin acetate in cardiac surgery: a double-blind, randomized study. *South Med J* 1994; 87:611-615

304. Terada N, Arai Y, Matsuta Y, Maekawa S, Okubo K, Ogura K, Matsuda N, Yonei A: Acute normovolemic hemodilution for radical prostatectomy: can it replace preoperative autologous blood transfusion? *Int J Urol* 2001; 8:149-152
305. Thomas D, Wareham K, Cohen D, Hutchings H: Autologous blood transfusion in total knee replacement surgery. *Br J Anaesth* 2001; 86:669-673
306. Tobias JD: Synthetic factor VIIa to treat dilutional coagulopathy during posterior spinal fusion in two children. *Anesthesiology* 2002; 96:1522-1525
307. Triulzi DJ, Gilmore GD, Ness PM, Baumgartner WA, Schultheis LW: Efficacy of autologous fresh whole blood or platelet-rich plasma in adult cardiac surgery. *Transfusion* 1995; 35:627-634
308. Troianos CA, Sypula RW, Lucas DM, D'Amico F, Mathie TB, Desai M, Pasqual RT, Pellegrini RV, Newfeld ML: The effect of prophylactic epsilon-aminocaproic acid on bleeding, transfusions, platelet function, and fibrinolysis during coronary artery bypass grafting. *Anesthesiology* 1999; 91:430-435
309. Trubel W, Gunen E, Wuppinger G, Tschernko E, Gunen-Frank A, Staudacher M, Polterauer P: Recovery of intraoperatively shed blood in aortoiliac surgery: comparison of cell washing with simple filtration. *Thorac Cardiovasc Surg* 1995; 43:165-170
310. Tsuji Y, Kambayashi J, Shiba E, Sakon M, Kawasaki T, Mori T: Effect of recombinant human erythropoietin on anaemia after gastrectomy: a pilot study. *Eur J Surg* 1995; 161:29-33
311. Uen WC, Chou YH, Liu CC, Lin SM, Chen TJ: Successful resection of sigmoid colon cancer in a patient with factor XI deficiency. *J Formos Med Assoc* 1998; 97:283-285
312. Urban MK, Beckman J, Gordon M, Urquhart B, Boachie-Adjei O: The efficacy of antifibrinolytics in the reduction of blood loss during complex adult reconstructive spine surgery. *Spine* 2001; 26:1152-1156
313. Veien M, Sorensen JV, Madsen F, Juelsgaard P: Tranexamic acid given intraoperatively reduces blood loss after total knee replacement: a randomized, controlled study. *Acta Anaesthesiol Scand* 2002; 46:1206-1211
314. Vichinsky EP, Haberkern CM, Neumayr L, Earles AN, Black D, Koshy M, Pegelow C, Abboud M, Ohene-Frempong K, Iyer RV: A comparison of conservative and aggressive transfusion regimens in the perioperative management of sickle cell disease. The Preoperative Transfusion in Sickle Cell Disease Study Group. *N Engl J Med* 1995; 333:206-213
315. Wajon P, Gibson J, Calcroft R, Hughes C, Thrift B: Intraoperative plateletpheresis and autologous platelet gel do not reduce chest tube drainage or allogeneic blood transfusion after reoperative coronary artery bypass graft. *Anesth Analg* 2001; 93:536-542
316. Walpoth B, Galliker B, Spirig P, Haeberli A, Rosenmund A, Althaus U, Nydegger UE: Use of epoetin alfa in autologous blood donation programs for patients scheduled for elective cardiac surgery. *Semin Hematol* 1996; 33(2 Suppl 2):75-76
317. Wang MJ, Chen TL, Huang CH, Chao A, Chu SH: Difference in efficacy of the Cell Saver in coronary bypass grafting surgery and cardiac valvular reoperations. *J Formos Med Assoc* 1994; 93:117-121
318. Waters JH, Lee JS, Klein E, O'Hara J, Zippe C, Potter PS: Preoperative autologous donation versus cell salvage in the avoidance of allogeneic transfusion in patients undergoing radical retropubic prostatectomy. *Anesth Analg* 2004 ;98:537-542
319. Waters JH, Lukauskiene E, Anderson ME: Intraoperative blood salvage during cesarean delivery in a patient with beta thalassemia intermedia. *Anesth Analg* 2003; 97:1808-1809
320. Waters JH, Sprung J: Errors during intraoperative cell salvage because of inappropriate wash solutions. *Anesth Analg* 2001; 93:1483-1485

321. Weisbach V, Skoda P, Rippel R, Lauer G, Glaser A, Zingsem J, Zimmermann R, Eckstein R: Oral or intravenous iron as an adjuvant to autologous blood donation in elective surgery: a randomized, controlled study. *Transfusion* 1999; 39:465-472
322. Whitten CW, Allison PM, Latson TW, Ivy R, Burkhardt D, Gulden RH, Cochran RP: Evaluation of laboratory coagulation and lytic parameters resulting from autologous whole blood transfusion during primary aortocoronary artery bypass grafting. *J Clin Anesth* 1996; 8:229-235
323. Wilhelmi M, Franke U, Cohnert T, Weber P, Kaukemuller J, Fischer S, Wahlers T, Haverich A: Coronary artery bypass grafting surgery without the routine application of blood products: Is it feasible? *Eur J Cardiothorac Surg* 2001; 19:657-661
324. Williams GD, Bratton SL, Riley EC, Ramamoorthy C: Coagulation tests during cardiopulmonary bypass correlate with blood loss in children undergoing cardiac surgery. *J Cardiothorac Vasc Anesth* 1999; 13:398-404
325. Wolowczyk L, Lewis DR, Nevin M, Smith FC, Baird RN, Lamont PM: The effect of acute normovolaemic haemodilution on blood transfusion requirements in abdominal aortic aneurysm repair. *Eur J Vasc Endovasc Surg* 2001; 22:361-364
326. Wolowczyk L, Nevin M, Smith FC, Baird RN, Lamont PM: Haemodilutional effect of standard fluid management limits the effectiveness of acute normovolaemic haemodilution in AAA surgery--results of a pilot trial. *Eur J Vasc Endovasc Surg* 2003; 26:405-411
327. Wong AY, Irwin MG, Hui TW, Fung SK, Fan ST, Ma ES: Desmopressin does not decrease blood loss and transfusion requirements in patients undergoing hepatectomy. *Can J Anaesth* 2002; 50:14-20
328. Wong JC, Torella F, Haynes SL, Dalrymple K, Mortimer AJ, McCollum CN; ATIS Investigators: Autologous versus allogeneic transfusion in aortic surgery: a multicenter randomized clinical trial. *Ann Surg* 2002; 235:145-151
329. Yamada T, Yamashita Y, Terai Y, Ueki M: Intraoperative blood salvage in abdominal uterine myomectomy. *Int J Gynaecol Obstet* 1997; 56:141-145
330. Yasuda H, Ohto H, Yamaguchi O, Sakuma S, Suzuki T, Mita M, Tsuneyama H, Uchikawa M: Three episodes of delayed hemolytic transfusion reactions due to multiple red cell antibodies, anti-Di, anti-Jk and anti-E. *Transfus Sci* 2000; 23:107-112
331. Yokomuro M, Ebine K, Shiroma K, Tamura S, Kumabe S, Ohtuki M, Suzuki H, Uchida S: Safety and efficacy of autologous platelet transfusion in cardiac surgery: comparison of cryopreservation, blood collection on the day before surgery, and blood collection during surgery. *Cryobiology* 1999; 38:236-242
332. Zabeeda D, Medalion B, Sverdlow M, Ezra S, Schachner A, Ezri T, Cohen AJ: Tranexamic acid reduces bleeding and the need for blood transfusion in primary myocardial revascularization. *Ann Thorac Surg* 2002; 74:733-738
333. Zupancic Salek S, Sokolic V, Viskovic T, Sanjug J, Simic M, Kastelan M: Successful use of recombinant factor VIIa for massive bleeding after caesarean section due to HELLP syndrome. *Acta Haematol* 2002; 108:162-163