

Supplementary Table S2 Differences between postinduction, intraoperative, and postoperative mean arterial pressure and ambulatory mean arterial pressure

Mean arterial pressure	Daytime, mean value		Daytime, lowest value		Nighttime, mean value		Nighttime, lowest value	
	Difference (perioperative - ambulatory), median (25th - 75th percentile), mmHg	Proportion (%) of patients with perioperative < ambulatory values	Difference (perioperative - ambulatory), median (25th - 75th percentile), mmHg	Proportion (%) of patients with perioperative < ambulatory values	Difference (perioperative - ambulatory), median (25th - 75th percentile), mmHg	Proportion (%) of patients with perioperative < ambulatory values	Difference (perioperative - ambulatory), median (25th - 75th percentile), mmHg	Proportion (%) of patients with perioperative < ambulatory values
Early postinduction, lowest value	-36 (-45 to -28)**	358/365 (98%)	-12 (-23 to -3)**	293/365 (80%)	-19 (-29 to -11)**	341/365 (93%)	-5 (-16 to 4)**	234/365 (64%)
Late postinduction, lowest value	-39 (-46 to -31)**	340/342 (99%)	-14 (-25 to -6)**	296/342 (87%)	-21 (-32 to -13)**	333/342 (97%)	-8 (-18 to 1)**	247/342 (72%)
Early intraoperative, lowest value	-36 (-44 to -26)**	366/368 (100%)	-11 (-22 to -1)**	278/368 (76%)	-19 (-28 to -11)**	333/368 (91%)	-6 (-14 to 4)**	244/368 (66%)
Late intraoperative, lowest value	-30 (-40 to -23)**	258/267 (97%)	-7 (-17 to 5)**	178/267 (67%)	-14 (-23 to -4)**	230/267 (86%)	0 (-9 to 8) ^{ns}	128/267 (48%)
Postoperative, lowest value	-25 (-37 to -14)**	295/326 (91%)	-2 (-13 to 12) ^{ns}	174/326 (53%)	-10 (-19 to 3)**	227/326 (70%)	+5 (-4 to 17)**	112/326 (34%)

^{ns} not significant

** p<0.001 Wilcoxon paired