

Tinnitus and Sleep Difficulties after Cochlear Implantation

Supplemental Digital Content 1

Robert H. Pierzycki,^{1,2} Mark Edmondson-Jones,^{1,2} Piers Dawes,³ Kevin J. Munro,^{3,4} David R. Moore,^{1,2,5} and Pádraig T. Kitterick^{1,2}

¹NIHR Nottingham Hearing Biomedical Research Unit, Ropewalk House, 113 The Ropewalk, Nottingham NG1 5DU, UK.

²Otology and Hearing Group, Division of Clinical Neuroscience, School of Medicine, University of Nottingham, Nottingham NG7 2UH, UK.

³School of Psychological Sciences, Ellen Wilkinson Building, University of Manchester, Oxford Road, Manchester M13 9PL, UK.

⁴Central Manchester University Hospitals NHS Foundation Trust, Manchester Academic Health Science Centre, Oxford Road, Manchester, M13 9WL, UK.

⁵Cincinnati Children's Hospital Medical Center and Department of Otolaryngology, University of Cincinnati College of Medicine, Cincinnati, OH 45229, USA.

GEOGRAPHICAL DISTRIBUTION OF PARTICIPANTS

Figure S1. Distribution of participants in the cochlear implant (CI) and Candidate groups across England and Wales based on the participants' local/unitary authority of residence. There were no assessments in Scotland and Northern Ireland at the time of the study.

