

TABLE E-1 Number of Radiolucent Lines and Radiolucent Line Scores on Anteroposterior and Axial Views by Glenoid Type

Glenoid Type <sup>23</sup>	Radiolucent Line Score <sup>24</sup> (points)								
	No.	Anteroposterior View				Axial View			
		Mean	Stand Dev.	Min.	Max.	Mean	Stand Dev.	Min.	Max.
A2	11	8.1	6.8	2	18	7.7	6.8	2	18
B1	16	7.4	6.5	2	18	7.3	6.5	2	18
B2	10	10.8	7.3	2	18	10.6	8.1	2	18
C	2	13	7.1	8	18	15.5	3.5	13	18

TABLE E-2 Influence of Cranial Migration of the Humeral Component on Clinical and Radiographic Outcomes

Migration	No.	Variable*	Mean	Stand. Dev.	Min.	Max.
None	12	Constant score (points)	63.3	13.2	45	81
		Pain (points)	11.6	2.9	5	15
		Flexion (°)	138.3	31	90	180
		Abduction (°)	126.7	34.9	70	180
		Strength (°)	6.5	1.9	4	10
		RLL AP (points)	7	6.7	2	18
		RLL axial (points)	6.9	6.8	2	18
Mild	10	Constant score (points)	65.4	19.1	28	86
		Pain (points)	12.6	3.1	6	15
		Flexion (°)	144	32	80	170
		Abduction (°)	142	39.1	70	170
		Strength (°)	6.6	5.6	0	15
		RLL AP (points)	6.7	6.3	2	2
		RLL axial (points)	6.5	6.3	18	18
Moderate	12	Constant score (points)	61.6	15.7	29	82
		Pain (points)	11.1	3.9	3	15
		Flexion (°)	132	37.3	40	170
		Abduction (°)	118.8	35.7	40	170
		Strength (°)	5.7	3.5	0	13
		RLL AP (points)	11.7	6.8	2	18
		RLL axial (points)	11.5	6.9	2	18
Severe	5	Constant score (points)	43.4	27.7	21	75
		Pain (points)	11	4.1	4	15
		Flexion (°)	98	44.9	60	160
		Abduction (°)	80	43	40	140
		Strength (°)	4	5.5	0	10
		RLL AP (points)	10.2	7.4	2	18
		RLL axial (points)	10.6	8.1	2	18

\*RLL AP = radiolucent line score on the anteroposterior view, and RLL axial = radiolucent line score on the axial view.