

TABLE E-1 Patient Demographics and Results of Treatment

Patient	Classification of Distal Arthrogyriposis	Diméglio Grade	Previous Treatment*	Number of Casts After Referral	Brace	Recurrence	Additional Treatment*	Ankle Dorsiflexion		Duration of Follow-up (yr)
								Right	Left	
1	DA2A	IV	NA	6	Dobbs	No	NA	10°	10°	2
2	DA2A	IV	NA	8	Semeda	No	NA	5°	10°	2
3	DA1A	IV	NA	8	Dobbs	No	NA	15°	10°	3
4	DA1A	IV	Casting	5	Dobbs	No	NA	5°	10°	3
5	DA1A	IV	Casting	8	Markell	Yes	(1) Repeat casting bilaterally. (2) Repeat Achilles tendon tenotomy on left.	20°	20°	2
6	DA1A	IV	NA	8	Dobbs	No	NA	15°	15°	2
7	DA1A	IV	NA	10	Semeda	Yes (twice)	(1) Repeat casting bilaterally. (2) Bilateral extensive soft-tissue-release operation.	0°	0°	2
8	DA4	IV	NA	6	Dobbs	No	NA	25°	25°	3
9	DA2A	IV	NA	7	Dobbs	No	NA	15°	20°	2
10	DA1A	IV	Casting	9	Markell	Yes	Repeat casting	20°	20°	4
11	DA5	IV	NA	6	Dobbs	No	NA	20°	20°	3
12	DA1A	II	Casting	2	Dobbs	No	NA	25°	25°	2

*NA = Not applicable.