

Appendix Table A1. Details of medical treatment used in each patient.

Case	GAT-IOP (mmHg)		Medical treatment			
	Before treatment	After treatment	Topical			IV 20% mannitol
			TDFC	Brimonidine	Latanoprost	
1	41	14	✓	✓	✓	once
2	52	10	✓	✓	✓	twice
3	48	23	✓	✓	✓	once
4	58	15	✓	✓	✓	once
5	48	14	✓	✓	✓	once
6	40	22	✓	✓	✓	once
7	38	10	✓	✓	✓	once
8	58	14	✓	✓	✓	once
9	45	20	✓	✓	✓	once
10	38	10	✓	✓	✓	none
11	58	10	✓	✓	✓	once
12	42	16	✓	✓	✓	once
13	64	10	✓	✓	✓	once
14	60	8	✓	✓	✓	twice
15	40	18	✓	✓	✓	once
16	60	12	✓	✓	✓	twice
17	58	16	✓	✓	✓	twice
18	56	15	✓	✓	✓	once
19	43	13	✓	✓	✓	once
20	60	12	✓	✓	✓	twice
21	48	24	✓	✓	✓	once
22	53	21	✓	✓	✓	once

Each topical medication was administered one drop at 5 minutes interval.

IV 20% mannitol at a dose of 300 ml were administered once or twice according to intraocular pressure (IOP) with at least 2 hours interval.

GAT = Goldmann applanation tonometry; TDFC = timolol/dorzolamide fixed combination; IV = intravenous