

Supplemental Table 1. Questionnaire Data by Sex and Age Group**For Ages < 20**

Female					
	Mean	SD	N	Median	IQR
BPI severity	5.88	1.72	143	6.00	2.50
BPI interference	6.07	2.29	146	6.43	3.71
DASS depression	16.48	11.97	143	16.00	20.00
DASS anxiety	12.78	10.16	144	10.00	16.00
DASS stress	19.73	10.80	143	20.00	14.00
PCS rumination	9.39	4.93	140	10.00	9.00
PCS magnification	5.56	3.39	140	6.00	5.00
PCS helplessness	13.50	6.56	139	15.00	9.00
PCS total	28.37	13.60	142	29.00	20.25
PSEQ total	24.48	13.27	145	23.00	18.00
Male					
	Mean	SD	N	Median	IQR
BPI severity	5.82	1.46	39	6.00	2.0
BPI interference	6.46	1.91	39	6.43	2.7
DASS depression	18.89	11.05	38	22.00	18.5
DASS anxiety	14.26	10.15	38	13.00	15.0
DASS stress	20.79	10.91	38	22.00	18.5
PCS rumination	10.08	4.63	39	11.00	6.0
PCS magnification	6.00	3.18	38	6.00	5.0
PCS helplessness	13.84	6.18	38	16.00	7.5
PCS total	29.67	12.94	39	31.00	16.0
PSEQ total	23.45	12.10	38	22.00	15.5

**For Ages 21-30,
Female**

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.03	1.63	565	6.00	2.25
BPI interference	6.88	2.02	570	7.14	2.71
DASS depression	18.41	12.74	557	16.00	20.00
DASS anxiety	13.76	10.46	556	14.00	16.00
DASS stress	21.40	10.95	557	22.00	18.00
PCS rumination	9.45	4.74	546	10.00	7.00
PCS magnification	5.67	3.48	549	5.00	5.00
PCS helplessness	14.14	6.20	547	14.00	9.00
PCS total	29.12	13.34	552	29.00	21.00
PSEQ total	21.37	12.47	567	19.00	17.00

Male

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.04	1.82	343	6.00	2.3
BPI interference	7.10	2.07	347	7.43	2.7
DASS depression	21.76	13.28	338	22.00	24.0
DASS anxiety	14.05	10.97	338	12.00	16.0
DASS stress	22.86	11.09	339	24.00	14.0
PCS rumination	10.65	4.81	325	12.00	8.0
PCS magnification	6.53	3.55	329	7.00	5.0
PCS helplessness	14.70	6.49	323	16.00	10.0
PCS total	31.72	13.80	329	34.00	22.0
PSEQ total	20.11	13.10	342	18.50	18.0

For Ages 31-40,**Female**

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.42	1.78	963	6.50	2.5
BPI interference	7.33	2.00	984	7.71	2.57
DASS depression	20.27	13.02	958	20.00	22.00
DASS anxiety	15.01	11.40	958	13.00	18.00
DASS stress	22.33	11.36	951	24.00	18.00
PCS rumination	9.67	4.71	931	10.00	8.00
PCS magnification	5.90	3.47	926	6.00	6.00
PCS helplessness	14.55	6.20	924	15.00	10.00
PCS total	29.97	13.18	936	31.00	21.00
PSEQ total	18.99	12.92	978	17.00	18.00

Male

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.39	1.76	800	6.50	2.5
BPI interference	7.28	1.95	819	7.57	2.7
DASS depression	22.69	12.49	791	24.00	22.0
DASS anxiety	14.51	10.99	784	12.00	17.5
DASS stress	23.41	10.81	780	24.00	18.0
PCS rumination	10.62	4.64	771	12.00	8.0
PCS magnification	6.49	3.49	771	6.00	5.0
PCS helplessness	14.86	6.19	751	16.00	10.0
PCS total	31.79	13.36	772	34.00	22.0
PSEQ total	18.65	13.10	806	16.00	19.0

For Ages 41-50,**Female**

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.46	1.78	1597	6.50	2.50
BPI interference	7.29	1.97	1639	7.57	2.71
DASS depression	20.80	13.04	1588	20.00	22.00
DASS anxiety	14.79	11.49	1583	12.00	18.00
DASS stress	21.62	11.50	1573	22.00	18.00
PCS rumination	9.71	5.03	1538	10.00	8.00
PCS magnification	5.92	3.64	1537	6.00	6.00
PCS helplessness	14.42	6.47	1524	15.00	10.00
PCS total	29.91	14.09	1552	31.00	23.00
PSEQ total	19.15	12.47	1615	18.00	18.00

Male

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.39	1.74	1284	6.50	2.5
BPI interference	7.27	1.95	1314	7.71	2.6
DASS depression	22.57	12.51	1267	22.00	22.0
DASS anxiety	14.64	11.08	1268	12.00	16.0
DASS stress	22.75	11.05	1260	24.00	18.0
PCS rumination	10.58	4.63	1230	12.00	8.0
PCS magnification	6.43	3.44	1231	7.00	5.0
PCS helplessness	14.56	6.21	1219	15.00	9.0
PCS total	31.44	13.10	1245	33.00	20.0
PSEQ total	18.68	12.76	1279	16.00	17.0

**For Ages 51-60,
Female**

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.54	1.80	1748	6.50	2.50
BPI interference	7.19	2.00	1795	7.57	2.57
DASS depression	20.43	12.91	1735	20.00	22.00
DASS anxiety	14.60	11.07	1734	12.00	16.00
DASS stress	20.84	11.62	1727	20.00	18.00
PCS rumination	9.52	5.02	1672	10.00	9.00
PCS magnification	5.83	3.62	1675	6.00	6.00
PCS helplessness	14.03	6.55	1654	15.00	10.00
PCS total	29.22	14.03	1696	30.00	22.75
PSEQ total	20.43	13.30	1776	19.00	18.00

Male

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.36	1.77	1344	6.50	2.25
BPI interference	7.15	1.98	1376	7.43	2.71
DASS depression	22.56	12.55	1323	24.00	22.00
DASS anxiety	14.70	10.94	1319	12.00	16.00
DASS stress	22.62	11.06	1315	22.00	18.00
PCS rumination	10.64	4.67	1277	12.00	8.00
PCS magnification	6.46	3.52	1280	7.00	5.00
PCS helplessness	14.67	6.29	1262	15.00	10.00
PCS total	31.60	13.29	1292	33.00	20.00
PSEQ total	19.63	12.67	1343	18.00	17.00

**For Ages 61-70,
Female**

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.40	1.81	1106	6.50	2.31
BPI interference	6.75	2.08	1154	7.14	2.71
DASS depression	17.55	12.72	1097	16.00	22.00
DASS anxiety	12.85	10.46	1096	10.00	16.00
DASS stress	18.31	11.62	1100	18.00	20.00
PCS rumination	9.21	5.06	1063	10.00	9.00
PCS magnification	5.41	3.58	1060	5.00	6.00
PCS helplessness	12.92	6.74	1035	13.00	11.00
PCS total	27.46	14.19	1069	28.00	23.00
PSEQ total	23.47	13.72	1123	23.00	20.00

Male

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.25	1.83	830	6.25	2.50
BPI interference	6.74	2.04	847	7.00	3.00
DASS depression	19.65	12.58	812	18.00	20.00
DASS anxiety	13.58	10.33	797	12.00	14.00
DASS stress	20.45	11.53	812	20.00	18.00
PCS rumination	10.07	4.87	804	11.00	8.00
PCS magnification	5.72	3.69	797	5.00	6.00
PCS helplessness	13.53	6.58	788	14.00	11.00
PCS total	29.10	14.00	809	29.00	23.00
PSEQ total	22.55	13.51	833	21.00	19.00

**For Ages 71-80,
Female**

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.68	1.96	787	6.75	2.75
BPI interference	6.68	2.17	843	6.86	2.86
DASS depression	17.21	12.09	784	16.00	20.00
DASS anxiety	12.77	9.98	775	10.00	14.00
DASS stress	17.86	11.61	776	16.00	18.00
PCS rumination	9.85	5.16	761	11.00	9.00
PCS magnification	5.54	3.76	753	5.00	6.00
PCS helplessness	13.54	7.03	733	14.00	12.00
PCS total	28.67	14.80	766	29.00	24.25
PSEQ total	24.24	14.19	807	23.00	21.00

Male

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.36	2.06	442	6.50	3.00
BPI interference	6.50	2.25	456	6.77	3.14
DASS depression	17.85	12.30	428	16.00	20.00
DASS anxiety	12.17	9.99	431	10.00	14.00
DASS stress	18.42	11.51	428	18.00	19.50
PCS rumination	10.14	4.97	430	11.00	9.00
PCS magnification	5.71	3.81	427	6.00	7.00
PCS helplessness	13.28	7.12	412	14.00	12.00
PCS total	28.83	14.72	436	30.00	24.75
PSEQ total	23.18	14.29	444	22.00	22.00

**For Ages 80+,
Female**

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.71	2.03	376	7.00	2.69
BPI interference	6.55	2.25	398	6.71	3.43
DASS depression	16.57	12.22	369	14.00	20.00
DASS anxiety	12.34	9.77	367	10.00	14.00
DASS stress	15.81	11.67	365	14.00	18.00
PCS rumination	9.77	5.27	363	10.00	10.00
PCS magnification	5.13	3.83	361	4.00	6.00
PCS helplessness	13.41	7.26	349	14.00	13.00
PCS total	28.23	15.26	358	30.00	27.00
PSEQ total	21.38	13.97	375	21.00	22.00

Male

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.33	2.08	152	6.50	3.50
BPI interference	6.48	2.20	161	6.71	3.15
DASS depression	18.46	12.98	153	16.00	24.00
DASS anxiety	11.86	10.14	152	10.00	12.00
DASS stress	17.91	12.48	152	18.00	22.00
PCS rumination	9.91	5.23	146	11.00	9.00
PCS magnification	5.46	3.69	146	6.00	6.00
PCS helplessness	13.22	7.25	142	14.00	11.00
PCS total	28.41	14.82	147	32.00	22.00
PSEQ total	24.64	15.65	154	22.50	24.25

Supplemental Table 2. Questionnaire Data by Work Status

Questionnaires	<u>Full time</u>					<u>Part-time</u>				
	Mean	SD	N	Median	IQR	Mean	SD	N	Median	IQR
BPI severity	5.65	1.87	1198	5.75	2.50	5.85	1.72	976	6.00	2.25
BPI interference	6.05	2.28	1223	6.29	3.14	6.43	2.05	1002	6.71	2.86
DASS depression	14.52	11.73	1190	12.00	18.00	16.68	12.30	966	14.00	20.00
DASS anxiety	9.41	9.06	1191	6.00	12.00	11.24	9.94	962	8.00	16.00
DASS stress	17.45	10.97	1188	16.00	18.00	18.93	11.41	966	18.00	18.00
PCS rumination	8.43	5.01	1156	9.00	9.00	8.66	4.85	940	9.00	8.00
PCS magnification	4.94	3.41	1161	5.00	5.00	5.21	3.34	942	5.00	5.00
PCS helplessness	11.84	6.49	1147	12.00	11.00	12.56	6.19	926	13.00	9.00
PCS total	25.11	13.74	1168	25.00	22.00	26.33	13.19	945	26.00	21.00
PSEQ total	27.64	13.67	1207	27.00	20.00	25.57	12.27	992	25.00	16.00

RetiredUnemployed due to pain

Questionnaires	Mean	SD	N	Median	IQR	Mean	SD	N	Median	IQR
BPI severity	6.45	1.93	3025	6.50	3.00	6.65	1.65	4278	6.75	2.25
BPI interference	6.71	2.15	3160	7.00	3.10	7.70	1.67	4380	8.00	2.29
DASS depression	17.93	12.53	3002	16.00	20.00	24.20	12.24	4210	26.00	20.00
DASS anxiety	12.96	10.25	2986	10.00	16.00	16.53	11.12	4208	16.00	19.50
DASS stress	18.38	11.77	2990	18.00	20.00	23.90	10.87	4181	24.00	16.00
PCS rumination	9.80	5.09	2941	11.00	8.00	10.72	4.59	4098	12.00	8.00
PCS magnification	5.55	3.70	2927	5.00	6.00	6.61	3.49	4085	7.00	5.00
PCS helplessness	13.36	6.93	2857	14.00	11.00	15.43	6.02	4038	16.00	9.00
PCS total	28.54	14.51	2961	29.00	24.00	32.62	12.96	4129	34.00	19.00
PSEQ total	22.86	13.93	3060	22.00	20.00	16.20	11.32	4315	15.00	16.00

Questionnaires	Unemployed not due to pain					Home duties				
	Mean	SD	N	Median	IQR	Mean	Std. Deviation	N	Median	IQR
BPI severity	6.51	1.79	671	6.50	2.50	6.54	1.85	1629	6.50	2.50
BPI interference	7.13	2.04	686	7.43	2.86	6.98	2.06	1692	7.29	2.71
DASS depression	23.29	12.69	653	24.00	20.00	19.45	12.66	1631	18.00	22.00
DASS anxiety	16.63	11.52	649	16.00	19.00	14.24	10.82	1631	12.00	16.00
DASS stress	23.51	11.18	646	24.00	18.00	20.67	11.65	1627	20.00	18.00
PCS rumination	10.49	4.64	618	11.00	8.00	9.86	4.97	1598	11.00	8.00
PCS magnification	6.54	3.54	619	7.00	5.00	5.90	3.68	1594	6.00	6.00
PCS helplessness	14.96	6.28	609	16.00	10.00	14.16	6.68	1574	15.00	11.00
PCS total	31.82	13.38	623	34.00	21.00	29.74	14.28	1617	31.00	23.00
PSEQ total	19.55	13.24	666	17.00	19.00	21.84	13.38	1665	20.00	19.00

Questionnaires	On leave because of pain					Studying				
	Mean	SD	N	Median	IQR	Mean	SD	N	Median	IQR
BPI severity	6.41	1.68	765	6.50	2.25	6.04	1.84	494	6.00	2.31
BPI interference	7.60	1.71	774	7.86	2.43	6.62	2.17	501	7.00	3.00
DASS depression	21.73	12.57	761	22.00	20.00	17.47	12.52	492	16.00	22.00
DASS anxiety	14.65	10.97	757	12.00	16.00	12.64	10.31	491	10.00	16.00
DASS stress	22.35	11.33	757	24.00	18.00	19.87	10.96	492	20.00	16.00
PCS rumination	10.02	4.85	732	11.00	8.00	8.93	4.80	487	9.00	8.00
PCS magnification	5.90	3.59	732	6.00	6.00	5.40	3.46	491	5.00	5.00
PCS helplessness	14.34	6.46	730	15.00	10.00	13.20	6.37	478	14.00	9.00
PCS total	30.06	13.76	741	32.00	21.00	27.42	13.41	493	27.00	20.00
PSEQ total	17.56	11.78	762	16.00	17.00	23.54	13.34	495	22.00	20.00

Questionnaires	Voluntary work					Retraining				
	Mean	SD	N	Median	IQR	Mean	SD	N	Median	IQR
BPI severity	6.11	1.70	365	6.00	2.25	6.10	1.98	105	6.00	2.88
BPI interference	6.38	2.06	371	6.43	3.14	6.92	2.16	109	7.43	3.29
DASS depression	16.77	12.33	362	14.00	20.00	19.88	12.74	104	18.00	24.00
DASS anxiety	12.18	10.44	358	10.00	14.50	13.81	11.14	103	10.00	16.00
DASS stress	18.94	11.16	361	18.00	18.00	20.49	10.83	103	22.00	20.00
PCS rumination	8.59	4.93	355	9.00	9.00	9.44	5.17	101	11.00	9.00
PCS magnification	4.89	3.51	354	4.00	5.00	5.58	3.69	101	6.00	7.00
PCS helplessness	11.80	6.97	348	11.50	11.00	13.12	6.78	99	14.00	11.00
PCS total	25.12	14.31	356	24.50	25.00	27.97	14.58	102	29.50	26.00
PSEQ total	27.49	13.16	367	28.00	19.00	21.73	12.56	108	21.50	17.75

Limited hours

Questionnaires	Mean	SD	N	Median	IQR
BPI severity	6.04	1.61	523	6.00	2.00
BPI interference	6.93	1.88	538	7.29	2.57
DASS depression	18.55	12.27	531	16.00	20.00
DASS anxiety	12.57	10.26	533	10.00	16.00
DASS stress	20.44	10.77	525	20.00	18.00
PCS rumination	9.21	4.68	515	10.00	7.00
PCS magnification	5.28	3.32	516	5.00	5.00
PCS helplessness	13.21	6.09	504	14.00	10.00
PCS total	27.54	12.94	518	28.00	21.00
PSEQ total	21.78	11.77	533	21.00	16.00

Supplemental Table 3. Questionnaire Data By Compensation Case

Questionnaires					
Compensation Case					
	Mean	SD	N	Median	IQR
BPI severity	6.53	1.66	1630	6.50	2.25
BPI interference	7.46	1.87	1661	7.86	2.43
DASS depression	23.64	12.67	1593	24.00	24.00
DASS anxiety	16.31	11.51	1585	14.00	20.00
DASS stress	24.35	11.12	1582	26.00	18.00
PCS rumination	10.57	4.73	1532	12.00	8.00
PCS magnification	6.38	3.60	1530	6.00	5.00
PCS helplessness	14.91	6.33	1519	16.00	10.00
PCS total	31.65	13.59	1544	33.00	21.75
PSEQ total	17.22	11.78	1637	15.00	16.00

Questionnaires					
Not Compensation Case					
	Mean	SD	N	Median	IQR
BPI severity	6.38	1.84	10426	6.50	2.50
BPI interference	6.97	2.07	10754	7.29	2.86
DASS depression	19.69	12.78	10343	18.00	22.00
DASS anxiety	13.75	10.73	10314	12.00	16.00
DASS stress	20.47	11.50	10294	20.00	18.00
PCS rumination	9.85	4.94	10074	11.00	8.00
PCS magnification	5.87	3.61	10061	6.00	6.00
PCS helplessness	13.96	6.57	9901	15.00	10.00
PCS total	29.53	13.96	10161	31.00	22.00
PSEQ total	21.20	13.45	10532	20.00	19.00

Supplemental Table 4. Questionnaire Data By Triggering Event

Questionnaires	Injury at home					Injury at work/school				
	Mean	SD	N	Median	IQR	Mean	SD	N	Median	IQR
BPI severity	6.49	1.76	850	6.50	2.50	6.46	1.69	2614	6.50	2.50
BPI interference	7.10	1.98	879	7.33	2.71	7.34	1.88	2688	7.71	2.66
DASS depression	20.67	12.51	842	20.00	20.00	22.77	12.63	2598	24.00	22.00
DASS anxiety	14.31	10.69	844	12.00	16.00	15.43	11.24	2585	14.00	18.00
DASS stress	21.08	11.28	838	22.00	18.00	23.24	10.98	2576	24.00	18.00
PCS rumination	10.39	4.79	830	12.00	8.00	10.26	4.78	2509	11.00	8.00
PCS magnification	6.08	3.65	826	6.00	6.00	6.25	3.53	2505	6.00	6.00
PCS helplessness	14.18	6.50	812	15.00	11.00	14.54	6.32	2478	15.00	10.00
PCS total	30.58	13.81	831	32.00	22.00	30.82	13.57	2530	32.50	21.00
PSEQ total	20.82	12.91	870	19.00	19.00	18.60	12.40	2642	17.00	17.00

Questionnaires	Injury in another setting					After surgery				
	Mean	SD	N	Median	IQR	Mean	SD	N	Median	IQR
BPI severity	6.36	1.82	909	6.25	2.75	6.28	1.91	1331	6.25	2.75
BPI interference	7.14	2.05	932	7.43	2.71	6.81	2.14	1380	7.14	2.71
DASS depression	20.15	13.00	888	20.00	24.00	18.78	12.64	1339	18.00	22.00
DASS anxiety	13.58	10.94	876	12.00	18.00	12.72	10.71	1336	10.00	16.00
DASS stress	20.91	11.68	879	20.00	18.00	19.53	11.84	1337	18.00	20.00
PCS rumination	9.80	4.85	858	10.00	8.00	9.77	5.01	1294	11.00	8.00
PCS magnification	5.87	3.59	859	6.00	6.00	5.62	3.65	1293	5.00	6.00
PCS helplessness	13.86	6.56	849	15.00	10.00	13.82	6.69	1278	15.00	10.00
PCS total	29.31	13.87	869	30.00	22.00	29.05	14.23	1310	30.00	23.00
PSEQ total	20.45	13.24	916	19.00	18.75	22.04	13.66	1345	21.00	20.00

Questionnaires	Motor vehicle crash					Related to cancer				
	Mean	SD	N	Median	IQR	Mean	Std. Deviation	N	Median	IQR
BPI severity	6.52	1.68	1161	6.50	2.50	5.93	1.99	212	6.00	2.50
BPI interference	7.29	1.90	1195	7.57	2.46	6.33	2.26	217	6.71	3.00
DASS depression	22.72	12.47	1160	24.00	22.00	16.31	12.15	211	14.00	20.00
DASS anxiety	16.25	11.22	1150	14.00	20.00	12.17	9.63	210	10.00	14.00
DASS stress	23.54	11.23	1150	24.00	18.00	17.92	11.86	212	16.00	17.50
PCS rumination	10.46	4.69	1128	11.00	8.00	8.37	4.87	199	8.00	8.00
PCS magnification	6.41	3.58	1125	6.00	5.00	4.74	3.56	199	4.00	5.00
PCS helplessness	14.89	6.22	1110	16.00	9.00	11.54	6.45	202	11.00	9.25
PCS total	31.59	13.36	1137	33.00	20.00	24.52	13.56	205	23.00	21.00
PSEQ total	18.88	12.62	1176	17.00	17.00	24.53	13.90	212	22.00	20.75

Questionnaires	Related to illness other than cancer					No obvious cause				
	Mean	SD	N	Median	IQR	Mean	SD	N	Median	IQR
BPI severity	6.34	1.84	1342	6.25	2.75	6.26	1.89	2171	6.25	2.50
BPI interference	7.01	2.02	1373	7.29	2.71	6.72	2.18	2254	7.00	3.00
DASS depression	19.13	12.68	1307	18.00	22.00	18.10	12.62	2175	16.00	20.00
DASS anxiety	14.19	10.50	1310	12.00	16.00	12.27	10.26	2163	10.00	14.00
DASS stress	19.88	11.27	1301	20.00	16.00	18.95	11.38	2164	18.00	18.00
PCS rumination	9.43	5.06	1285	10.00	9.00	9.74	4.95	2120	10.00	8.00
PCS magnification	5.67	3.55	1273	5.00	5.00	5.74	3.56	2139	6.00	5.00
PCS helplessness	13.77	6.61	1250	14.00	10.00	13.80	6.54	2081	15.00	10.00
PCS total	28.70	14.03	1286	30.00	22.00	29.13	13.86	2142	30.00	22.00
PSEQ total	20.73	12.94	1331	19.00	18.00	22.47	13.98	2224	21.00	20.00

Questionnaires	Other				
	Mean	SD	N	Median	IQR
BPI severity	6.54	1.89	1530	6.50	2.75
BPI interference	6.98	2.11	1584	7.29	2.86
DASS depression	19.51	13.15	1509	18.00	24.00
DASS anxiety	13.97	10.92	1520	12.00	16.00
DASS stress	20.47	11.72	1511	20.00	18.00
PCS rumination	9.99	4.91	1491	11.00	8.00
PCS magnification	5.96	3.70	1478	6.00	6.00
PCS helplessness	14.08	6.72	1465	15.00	11.00
PCS total	29.92	14.15	1505	31.00	24.00
PSEQ total	21.30	13.71	1544	20.00	18.00

Supplemental Table 5. Questionnaire Data By Missing Data

	Number missing	Percentage missing
BPIWorst	392	2.9%
BPILeast	517	3.9%
BPIAverage	464	3.5%
BPINow	463	3.5%
BPIActivity	432	3.2%
BPIMood	448	3.4%
BPIWalking	449	3.4%
BPIWork	538	4.0%
BPIRelations	570	4.3%
BPISleep	463	3.5%
BPIEnjoyment	537	4.0%
DASSQ1	946	7.1%
DASSQ2	822	6.2%
DASSQ3	972	7.3%
DASSQ4	894	6.7%
DASSQ5	776	5.8%
DASSQ6	831	6.2%
DASSQ7	839	6.3%
DASSQ8	948	7.1%
DASSQ9	807	6.0%
DASSQ10	791	5.9%
DASSQ11	801	6.0%
DASSQ12	764	5.7%
DASSQ13	886	6.6%
DASSQ14	1030	7.7%
DASSQ15	847	6.3%
DASSQ16	818	6.1%
DASSQ17	823	6.2%
DASSQ18	909	6.8%
DASSQ19	929	7.0%

Supplemental Table 5. Questionnaire Data By Missing Data

DASSQ20	851	6.4%
DASSQ21	877	6.6%
PCSQ1	861	6.5%
PCSQ2	908	6.8%
PCSQ3	891	6.7%
PCSQ4	962	7.2%
PCSQ5	939	7.0%
PCSQ6	928	7.0%
PCSQ7	1007	7.5%
PCSQ8	943	7.1%
PCSQ9	941	7.1%
PCSQ10	931	7.0%
PCSQ11	953	7.1%
PCSQ12	952	7.1%
PCSQ13	975	7.3%
PSEQQ1	540	4.0%
PSEQQ2	547	4.1%
PSEQQ3	555	4.2%
PSEQQ4	532	4.0%
PSEQQ5	578	4.3%
PSEQQ6	564	4.2%
PSEQQ7	559	4.2%
PSEQQ8	571	4.3%
PSEQQ9	534	4.0%
PSEQQ10	587	4.4%