

Table E1. Best-corrected visual acuity before and after treatment with intravenous erythropoietin and measurements of the peripapillary retinal nerve fiber layer (PPRNFL)

Case No.	Right Mean PPNFL Thickness (μm)	Left Mean PPNFL Thickness (μm)	Follow-up (months)	Initial VA		Final VA (at least 3 month)	
				OD	OS	OD	OS
1.	50.25	51.00	23	NLP	NLP	20/20	20/20
2.	NA	NA	18	HM	HM	20/40	20/30
3.	55.00	64.75	5	NLP	HM	HM	20/400
4.	NA	NA	8	NLP	HM	20/25	20/22.5
5.	NA	NA	5	LP	HM	CF	20/22.5
6.	NA	NA	3	NLP	NLP	CF	CF
7.	NA	NA	4	NLP	NLP	NLP	NLP
8.	NA	NA	3	NLP	LP	20/32	20/25
9.	77.75	61.50	6	HM	HM	CF	CF
10.	NA	NA	5	HM	LP	20/120	CF
11.	106.00	115.50	4	NLP	NLP	20/32	HM
12.	NA	NA	15	20/400	20/400	20/300	20/200
13.	NA	NA	3	CF	20/80	CF	20/50
14.	NA	NA	7	NLP	NLP	20/32	20/40
15.	122.75	78.00	7	HM	HM	20/160	HM
16.	NA	NA	5	NLP	NLP	20/32	20/25